IHES/P/12/21, IPHT-t12/057

arXiv:1208.1255v2 [hep-th] 7 Sep 2012

One-loop four-graviton amplitudes in $\mathcal{N} = 4$ supergravity models

Piotr Tourkine[a] and **Pierre Vanhove**[a,b]

[a] *Institut de Physique Théorique, CEA/Saclay, F-91191 Gif-sur-Yvette, France*

[b] *IHES, Le Bois-Marie, 35 route de Chartres, F-91440 Bures-sur-Yvette, France*

email: `piotr.tourkine,pierre.vanhove@cea.fr`

Abstract

We evaluate with great details one-loop four-graviton field theory amplitudes in pure $\mathcal{N} = 4$ $D = 4$ supergravity. The expressions are obtained by taking the field theory limit of $(4,0)$ and $(2,2)$ space-time supersymmetric string theory models. For each model we extract the contributions of the spin 1 and spin 2 $\mathcal{N} = 4$ supermultiplets running in the loop. We show that all the string based constructions with $(4,0)$ symmetry lead to the same answer as the one derived by unitarity and double-copy method in field theory. But the limit of the string theory $(2,2)$ models leads to a different answer, that we attribute to the different nature of the vector multiplet couplings in the various string theory models.

I. INTRODUCTION

The role of supersymmetry in perturbative supergravity still leaves room to surprises. The construction of candidate counter-terms for ultraviolet (UV) divergences in extended four dimensional supergravity theories does not forbid some particular amplitudes to have an improved UV behaviour. For instance, the four-graviton three-loop amplitude in $\mathcal{N} = 4$ supergravity turns out to be UV finite [1, 2], despite the construction of a candidate counter-term [3]. (Some early discussion of the three-loop divergence in $\mathcal{N} = 4$ has appeared in [4], and recent alternative arguments have been given in [5].)

The UV behaviour of extended supergravity theories is constrained in string theory by non-renormalisation theorems that give rise in the field theory limit to supersymmetric protections for potential counter-terms. In maximal $\mathcal{N} = 8$ supergravity, the absence of divergences till six loops in four dimensions [6–8] is indeed a consequence of the supersymmetric protections for $\frac{1}{2}$-, $\frac{1}{4}$- and $\frac{1}{8}$-BPS operators in string [9, 10] or field theory [11, 12]. In $\mathcal{N} = 4$ supergravity, it was shown recently [2] that the absence of three-loop divergence in the four-graviton amplitude in four dimensions is a consequence of the protection of the $\frac{1}{2}$-BPS R^4 coupling from perturbative quantum corrections beyond one-loop in heterotic models. We refer to [13–15] for a discussion of the non-renormalisation theorems in heterotic string.

Maximal $\mathcal{N} = 8$ supergravity is unique in any dimension, and corresponds to the massless sector of type II string theory compactified on a torus. Duality symmetries relate different phases of the theory and strongly constrain its UV behaviour [10, 12, 16–19].

To the contrary $\mathcal{N} = 4$ supergravity (coupled to vector multiplets) is not unique and can be obtained in the low-energy limit of $(4, 0)$ string theory models—with all the space-time supersymmetries coming from the world-sheet left-moving sector—or $(2, 2)$ string theory models—with the space-time supersymmetries originating both from the world-sheet left-moving and right-moving sectors. The two constructions will give rise to different low-energy supergravity theories with a different identification of the moduli. We will see that as a consequence, the amplitudes differ in the two theories.

In this work we analyze the properties of the four-graviton amplitude at one-loop in $\mathcal{N} = 4$ supergravity. We write the genus one string theory amplitude in different models and extract its field theory limit. We consider three classes of four dimensional string models. The first class, on which was based the analysis in [2], are heterotic string models. They have $(4,0)$ supersymmetry and $4 \leq n_v \leq 22$ vector multiplets. The models of the second class also carry $(4,0)$ supersymmetry, they are type II asymmetric orbifolds. We will study a model with $n_v = 6$ vector multiplets as well as the Dabholkar-Harvey models with $n_v = 0$ vector multiplets [20]. The third class is composed of type II symmetric orbifolds with $(2,2)$ supersymmetry. For a given number of vector multiplets, the $(4,0)$ models are related to one another by strong-weak S-duality and related to $(2,2)$ models by U-duality [21, 22]. Several tests of the duality relations between orbifold models have been given in [23].

Our string theory constructions contain generically vector multiplets, however we will be able to isolate the contributions of the $\mathcal{N} = 4$ spin 1 and spin 2 supermultiplets to the field theory amplitude. We shall indeed show that the amplitudes extracted from the $(4,0)$ string models match the one obtained from unitarity in [24–29] and treating gravity as double-copy of gauge theory in [30]. However, the amplitudes in the $(2,2)$ construction take a different form. This is expected from the string theory duality relations, and it would be interesting to reproduce them from the double-copy construction of two set of $\mathcal{N} = 2$ super-Yang-Mills amplitudes.

The paper is organized as follows. For each model we evaluate the one-loop four-graviton string theory amplitude in section II. In section III we compare the expressions that we obtained and check that they are compatible with our expectations from string dualities. We then extract and evaluate the field theory limit in the regime $\alpha' \to 0$ of those string amplitudes in section IV. This gives us the field theory four-graviton one-loop amplitudes for pure $\mathcal{N} = 4$ supergravity. Section V contains our conclusion. Finally, the appendices A and B contain details about our conventions and the properties of the CFT building blocks of our string theory models.

II. ONE-LOOP STRING THEORY AMPLITUDES IN $(4,0)$ AND $(2,2)$ MODELS

In this section, we compute one-loop four-graviton amplitudes in four dimensional $\mathcal{N} = 4$ $(4,0)$ and $(2,2)$ string theory models. Their massless spectrum contains an $\mathcal{N} = 4$ supergravity multiplet coupled to n_v $\mathcal{N} = 4$ vector multiplets. Since the heterotic string is made of the tensor product of a left moving superstring by a right moving bosonic string, it only gives rise to $(4,0)$ models. However, type II compactifications provide the freedom to build $(4,0)$ and $(2,2)$ models [31].

A. Heterotic CHL models

We evaluate the one-loop four-graviton amplitudes in heterotic string CHL models in four dimensions [32–34]. Their low-energy limit are $(4,0)$ supergravity models with $4 \leq n_v \leq 22$ vector supermultiplets matter fields. We first comment on the moduli space of the model, and then write the string theory one-loop amplitude. Then, we compute the CHL partition function. This allows us to extract the massless states contribution to the integrand of the field theory limit.

These models have the following moduli space

$$\Gamma\backslash SU(1,1)/U(1) \times SO(6,n_v;\mathbb{Z})\backslash SO(6,n_v)/SO(6) \times SO(n_v)\,, \tag{II.1}$$

where n_v is number of vector multiplets, and Γ is a discrete subgroup of $SL(2,\mathbb{Z})$. For instance, $\Gamma = SL(2,\mathbb{Z})$ for $n_v = 22$ and $\Gamma = \Gamma_1(N)$ for the $\mathbb{Z}_N$ CHL $(4,0)$ orbifold. (We refer to appendix A 3 for a definition of the congruence subgroups of $SL(2,\mathbb{Z})$.) The scalar manifold $SU(1,1)/U(1)$ is parametrized by the axion-dilaton in the $\mathcal{N} = 4$ gravity supermultiplet.

The generic structure of the amplitude has been described in [2]. We will use the same notations and conventions. The four-graviton amplitude takes the following form

$$\mathcal{M}^{(n_v)}_{(4,0)het} = \mathcal{N}\left(\frac{\pi}{2}\right)^4 t_8F^4 \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{\frac{D-6}{2}}} \int_{\mathcal{T}} \prod_{1\leq i<j\leq 4} \frac{d^2\nu_i}{\tau_2}\, e^{\mathcal{Q}}\, \mathcal{Z}^{(n_v)}_{(4,0)het} \bar{\mathcal{W}}^B\,, \tag{II.2}$$

where $D = 10 - d$, and $\mathcal{N}$ is the normalization constant of the amplitude. The domains of integration are $\mathcal{F} = \{\tau = \tau_1 + i\tau_2; |\tau_1| \leq \frac{1}{2}, |\tau|^2 \geq 1, \tau_2 > 0\}$ and $\mathcal{T} := \{\nu = \nu_1 + i\nu_2; |\nu_1| \leq \frac{1}{2}, 0 \leq \nu_2 \leq \tau_2\}$. Then,

$$\bar{\mathcal{W}}^B := \frac{\langle \prod_{j=1}^4 \tilde{\epsilon}^j \cdot \bar{\partial}X(z_j)e^{ik_j\cdot X(z_j)}\rangle}{(2\alpha')^4\langle \prod_{j=1}^4 e^{ik_j\cdot X(z_j)}\rangle}\,, \tag{II.3}$$

is the kinematical factor coming from Wick contracting the bosonic vertex operators, and the plane-wave part is given by $\langle \prod_{j=1}^4 e^{ik_j\cdot X(z_j)}\rangle = \exp(\mathcal{Q})$ with

$$\mathcal{Q} = \sum_{1\leq i<j\leq 4} 2\alpha' k_i \cdot k_j \mathcal{P}(\nu_{ij})\,. \tag{II.4}$$

Using the result of [35] (but with our normalizations), we explicitly write

$$\mathcal{W}^B = \prod_{r=1}^4 \epsilon_r\cdot\mathcal{Q}_r + \frac{1}{2\alpha'}\left(\tilde{\epsilon}_1\cdot\mathcal{Q}_1\,\tilde{\epsilon}_2\cdot\mathcal{Q}_2\,\tilde{\epsilon}_3\cdot\tilde{\epsilon}_4\mathcal{T}(\nu_{34}) + perms\right) + \frac{1}{4\alpha'^2}\left(\tilde{\epsilon}_1\cdot\tilde{\epsilon}_2\,\tilde{\epsilon}_3\cdot\tilde{\epsilon}_4\,\mathcal{T}(\nu_{12})\mathcal{T}(\nu_{34}) + perms\right), \tag{II.5}$$

where we have introduced

$$\mathcal{Q}_I^\mu := \sum_{r=1}^4 k_r^\mu \, \partial\mathcal{P}(z_{Ir}|\tau); \qquad \mathcal{T}(\nu) := \partial_\nu^2 \mathcal{P}(\nu|\tau)\,. \tag{II.6}$$

and $\mathcal{P}(z)$ is the genus one bosonic propagator. We refer to appendix A 2 for definitions and conventions.

The CHL models studied in this work are asymmetric $\mathbb{Z}_N$ orbifolds of the bosonic sector (in our case the right moving sector) of the heterotic string compactified on $T^5 \times S^1$. Geometrically, the orbifold rotates N groups of ℓ bosonic fields $\bar{X}^a$ belonging either to the internal T^{16} either to the T^5 and act as an order N shift on the S^1. Precisely if we take a boson X^a of the $(p+1)$-th group ($p = 0, \dots, N-1$) of ℓ bosons we have $a \in \{p\ell, p\ell+1, \dots, p\ell+(\ell-1)\}$ and for twists $g/2, h/2 \in \{0, 1/N, \dots, 1(N-1)/N\}$ we get

$$\begin{aligned}\bar{X}^a(z+\tau) &= e^{i\pi g p/N} \bar{X}^a(z)\,,\\ \bar{X}^a(z+1) &= e^{i\pi h p/N} \bar{X}^a(z)\,.\end{aligned} \tag{II.7}$$

We will consider models with $(N, n_v, \ell) \in \{(1,22,16), (2,14,8), (3,10,6), (5,6,4), (7,4,3)\}$. It is in principle possible to build models with $(N, n_v, \ell) = (11,2,2)$ and $(N, n_v, \ell) = (23,0,1)$ and thus decouple totally the matter fields, but it is then required to compactify the theory on a seven- and eight-dimensional torus respectively. We will not comment about it further, since anyway we have a type II superstring with $(4,0)$ supersymmetry that already has $n_v = 0$ described in section II B 2. The partition function of the right moving CFT writes

$$\mathcal{Z}^{(n_v)}_{(4,0)het}(\tau) = \frac{1}{|G|} \sum_{(g,h)} \mathcal{Z}^{h,g}_{(4,0)het}(\tau)\,, \tag{II.8}$$

where $|G|$ is the order of the orbifold group *i.e.* $|G| = N$. The twisted conformal blocks $\mathcal{Z}^{g,h}_{het}$ are a product of the oscillator part and the zero mode part

$$\mathcal{Z}^{h,g}_{(4,0)het} = \mathcal{Z}^{h,g}_{osc} \times \mathcal{Z}^{h,g}_{latt}\,. \tag{II.9}$$

In the field theory limit only the massless states from the sector $h = 0$ will contribute and we are left with:

$$\mathcal{Z}^{(n_v)}_{(4,0)het}(\tau) \to \frac{1}{N}\,\mathcal{Z}^{0,0}_{(4,0)het}(\tau) + \frac{1}{N} \sum_{\{g\}} \mathcal{Z}^{0,g}_{(4,0)het}(\tau)\,. \tag{II.10}$$

The untwisted partition function ($g = h = 0$) with generic diagonal Wilson lines A, as required by modular invariance, is

$$\mathcal{Z}^{0,0}_{(4,0)het}(\tau) := \frac{\Gamma_{(6,24)}(G,A)}{\bar\eta^{24}(\bar\tau)}\,, \tag{II.11}$$

where $\Gamma_{(6,24)}(G,A)$ is the lattice sum for the Narain lattice $\Gamma^{(5,5)} \oplus \Gamma^{(1,1)} \oplus \Gamma_{E_8\times E_8}$ with Wilson lines [36]. It drops out in the field theory limit where the radii of compactification $R \sim \sqrt{\alpha'}$ are sent to zero and we are left with the part coming from the oscillators

$$\mathcal{Z}^{0,0}_{(4,0)het}(\tau) \to \frac{1}{\bar\eta^{24}(\bar\tau)}\,. \tag{II.12}$$

At a generic point in the moduli space, the 480 gauge bosons of the adjoint representation of $E_8 \times E_8$ get masses due to Wilson lines, and only the ℓ gauge bosons of the $U(1)^\ell$ group left invariant by the orbifold action [37, 38] stay in the matter massless spectrum.

The oscillator part is computed to be

$$\mathcal{Z}^{h,g}_{osc} = \sum_{\{g,h\}} \prod_{p=0}^{N-1} \left(\mathcal{Z}_X^{h\times p, g\times p}\right)^{\ell}\,, \tag{II.13}$$

where the twisted bosonic chiral blocks $\mathcal{Z}^{h,g}_X$ are given in appendix A. For $h = 0$, $\mathcal{Z}^{0,g}_{osc}$ is independent of g when N is prime and it can be computed explicitly. It is the inverse of the unique cusp form $f_k(\tau) = \eta(\tau)^{k+2}\eta(N\tau)^{k+2}$ for $\Gamma_1(N)$ of modular weight $\ell = k+2 = 24/(N+1)$ with $n_v = 2\ell - 2$ as determined in [37, 38]. Then (II.10) writes

$$\mathcal{Z}_{(4,0)het} \to \frac{1}{N}\left(\frac{1}{\bar\eta(\bar\tau)^{24}} + \frac{N-1}{f_k(\bar\tau)}\right)\,. \tag{II.14}$$

To conclude this section, we write the part of the integrand of (II.2) that will contribute in the field theory limit. When $\alpha' \to 0$, the region of the fundamental domain of integration $\mathcal{F}$ of interest is the large τ_2 region, such that $t = \alpha'\tau_2$ stays constant. Then, the objects that we have introduced admit an expansion in the variable $q = e^{2i\pi\tau} \to 0$. We find

$$\mathcal{Z}_{(4,0)het} \to \frac{1}{\bar q} + 2 + n_v + O(\bar q)\,. \tag{II.15}$$

Putting everything together and using the expansions given in (A.16), we get for the integrand in (II.2)

$$\mathcal{Z}_{(4,0)het}\mathcal{W}^B e^{\mathcal{Q}} \to e^{\pi\alpha'\tau_2 Q}\left(\left(\mathcal{W}^B e^{\mathcal{Q}}\right)|_{\bar q} + (n_v+2)\left(\mathcal{W}^B e^{\mathcal{Q}}\right)|_{\bar q^0} + O(\bar q)\right)\,. \tag{II.16}$$

The order $\bar{q}$ coefficients arise because of the presence of the $1/\bar{q}$ chiral tachyonic pole in the non-supersymmetric sector of the theory. Since the integral over τ_1 of $\bar{q}^{-1}\left(\mathcal{W}^B e^{\mathcal{Q}}\right)|_{\bar{q}^0}$ vanishes, as a consequence of the level matching condition, we did not write it. We introduce $\mathcal{A}$, the massless sector contribution to the field theory limit of the amplitude at the leading order in α', for later use in sections III and IV

$$\mathcal{A}^{(n_v)}_{(4,0)het} = \frac{1}{2}\left(\frac{\pi}{2}\right)^4 t_8F^4 \left(\bar{\mathcal{W}}^B|_{\bar{q}}(1+\alpha'\delta\mathcal{Q}) + \bar{\mathcal{W}}^B|_{\bar{q}^0}\mathcal{Q}|_{\bar{q}} + (n_v+2)\bar{\mathcal{W}}^B|_{\bar{q}^0}\right)\,, \tag{II.17}$$

where we have made used of the notations for the $\bar{q}$-expansion

$$\begin{aligned}\bar{\mathcal{W}}^B &= \bar{\mathcal{W}}^B|_{q^0} + \bar{q}\,\bar{\mathcal{W}}^B|_q + O(\bar{q}^2)\,,\\ \mathcal{Q} &= -\pi\,\alpha'\tau_2\,Q + \alpha'\delta Q + q\,\mathcal{Q}|_q + \bar{q}\,\mathcal{Q}|_{\bar{q}} + O(|q|^2)\,.\end{aligned} \tag{II.18}$$

B. Type II asymmetric orbifold

In this section we consider type II string on two different kinds of asymmetric orbifolds. They lead to $(4,0)$ models with a moduli space given in (II.1), where the axion-dilaton parametrizes the $SU(1,1)/U(1)$ factor. The first one is a $\mathbb{Z}_2$ orbifold with $n_v = 6$. The other ones are the Dabholkar-Harvey models [20, 39], they have with $n_v = 0$.

First, we give a general formula for the treatment of those asymmetric orbifolds. Then we study in detail the partition functions of two models, and extract the massless states contribution to the integrand of the field theory limit of the amplitudes. A generic expression for the scattering amplitude of four gravitons at one-loop in type IIA and IIB superstring is:

$$\begin{aligned}\mathcal{M}^{(n_v)}_{(4,0)II} = \mathcal{N}\int_{\mathcal{F}}\frac{d^2\tau}{\tau_2^{\frac{D-6}{2}}}\int_{\mathcal{T}}\prod_{1\leq i<j\leq 4}\frac{d^2\nu_i}{\tau_2}\,e^{\mathcal{Q}}\,\times\\ \times\frac{1}{2}\sum_{a,b=0,1}(-1)^{a+b+ab}\mathcal{Z}_{a,b}\mathcal{W}_{a,b}\,\times\\ \times\frac{1}{2|G|}\sum_{\substack{\bar{a},\bar{b}=0,1\\ g,h}}(-1)^{\bar{a}+\bar{b}+\mu\bar{a}\bar{b}}(-1)^{C(\bar{a},\bar{b},g,h)}\,\bar{\mathcal{Z}}^{h,g}_{\bar{a},\bar{b}}\tilde{\mathcal{W}}_{\bar{a},\bar{b}}\,,\end{aligned} \tag{II.19}$$

where $\mathcal{N}$ is the same normalization factor as for the heterotic string amplitude and $C(\bar{a},\bar{b},g,h)$ is a model dependent phase factor determined by modular invariance and discussed below. We have introduced the chiral partition functions in the (a,b)-spin structure

$$\mathcal{Z}_{a,b} = \frac{\theta\left[{a \atop b}\right](0|\tau)^4}{\eta(\tau)^{12}}\,;\qquad \mathcal{Z}_{1,1} = 0\,. \tag{II.20}$$

The value of μ determines the chirality of the theory: $\mu=0$ for type IIA and $\mu=1$ for type IIB. The partition function in a twisted sector (h,g) of the orbifold is $\bar{\mathcal{Z}}^{h,g}_{\bar a,\bar b}$. Notice that the four dimensional fermions are not twisted, so the vanishing of their partition function in the $(a,b)=(1,1)$ sector implies the vanishing of $\bar{\mathcal{Z}}^{h,g}_{1,1}=0$ for all g,h. This is fully consistent with the fact that this amplitude does not receive any contributions from the odd/odd, odd/even or even/odd spin structure by lack of fermionic zero modes. We use the holomorphic factorization of the $(0,0)$-ghost picture graviton vector operators as

$$V^{(0,0)}=\int d^2z\ :\epsilon^{(i)}\cdot V(z)\,\tilde\epsilon^{(i)}\cdot\bar V(\bar z)\,e^{ik\cdot X}:\,, \tag{II.21}$$

with

$$\epsilon^{(i)}\cdot V(z)=\epsilon^{(i)}\cdot\partial X-i\frac{F^{(i)}_{\mu\nu}}{2}:\psi^\mu\psi^\nu:;\qquad \tilde\epsilon^{(i)}\cdot\bar V(\bar z)=\tilde\epsilon^{(i)}\cdot\bar\partial X+i\frac{\tilde F^{(i)}_{\mu\nu}}{2}:\bar\psi^\mu\bar\psi^\nu:\,, \tag{II.22}$$

where we have introduced the field strengths $F^{(i)}_{\mu\nu}=\epsilon^{(i)}_\mu k_{i\,\nu}-\epsilon^{(i)}_\nu k_{i\,\mu}$ and $\tilde F^{(i)}_{\mu\nu}=\tilde\epsilon^{(i)}_\mu k_{i\,\nu}-\tilde\epsilon^{(i)}_\nu k_{i\,\mu}$.

The correlators of the vertex operators in the (a,b)-spin structure are given by $\mathcal{W}_{a,b}$ and $\bar{\mathcal{W}}_{\bar a,\bar b}$ defined by

$$\mathcal{W}_{a,b}=\frac{\langle\prod_{j=1}^4\epsilon^{(j)}\cdot V(z_j)\,e^{ik_j\cdot X(z_j)}\rangle_{a,b}}{(2\alpha')^4\langle\prod_{j=1}^4 e^{ik_j\cdot X(z_j)}\rangle}\,,\qquad \bar{\mathcal{W}}_{\bar a,\bar b}=\frac{\langle\prod_{j=1}^4\tilde\epsilon^{(j)}\cdot\bar V(\bar z_j)\,e^{ik_j\cdot X(z_j)}\rangle_{\bar a,\bar b}}{(2\alpha')^4\langle\prod_{j=1}^4 e^{ik_j\cdot X(\bar z_j)}\rangle}\,, \tag{II.23}$$

We decompose the $\mathcal{W}_{a,b}$ in a part that depends on the spin structure (a,b), denoted $\mathcal{W}^F_{a,b}$, and a part independent of the spin structure $\mathcal{W}^B$

$$\mathcal{W}_{a,b}=\mathcal{W}^F_{a,b}+\mathcal{W}^B\,, \tag{II.24}$$

this last contribution is identical to the one given in (II.3). The spin structure dependent part is given by the following fermionic Wick's contractions:

$$\mathcal{W}^F_{a,b}=\mathcal{S}_{4;a,b}+\mathcal{S}_{2;a,b}\,, \tag{II.25}$$

where $\mathcal{S}_{n;a,b}$ arise from Wick contracting n pairs of world-sheet fermions. Note that the contractions involving three pairs of fermion turns out to vanish in all the type II models by symmetry. Introducing the notation $\sum_{\{(i,\cdots),(j,\cdots)\}=\{1,2,3,4\}}\cdots$ for the sum over the ordered partitions of $\{1,2,3,4\}$ into two sets. In the sum, the partitions $\{(1,2,3),1\}$ and $\{(1,3,2),1\}$ are considered to be different. In that manner, the two terms in (II.25) can be written explicitly:

$$\begin{aligned}\mathcal{S}_{4;a,b} &= \frac{1}{2^{10}} \sum_{\{(i,j),(k,l)\}=\{1,2,3,4\}} S_{a,b}(z_{ij})S_{a,b}(z_{ji})S_{a,b}(z_{kl})S_{a,b}(z_{lk})\,\mathrm{tr}(F^{(i)}F^{(j)})\,\mathrm{tr}(F^{(k)}F^{(l)})\\ &\quad - \frac{1}{2^8} \sum_{\{(i,j,k,l)\}=\{1,2,3,4\}} S_{a,b}(z_{ij})S_{a,b}(z_{jk})S_{a,b}(z_{kl})S_{a,b}(z_{li})\,\mathrm{tr}(F^{(i)}F^{(j)}F^{(k)}F^{(l)}) \qquad \text{(II.26)}\\ \mathcal{S}_{2;a,b} &= -\frac{1}{2^5} \sum_{\{(i,j),(k,l)\}=\{1,2,3,4\}} S_{a,b}(z_{ij})S_{a,b}(z_{ji})\,\mathrm{tr}(F^{(i)}F^{(j)})\,(\epsilon^{(k)}\cdot\mathcal{Q}_k\,\epsilon^{(l)}\cdot\mathcal{Q}_l + \frac{1}{2\alpha'}\epsilon^{(k)}\cdot\epsilon^{(l)}\,\mathcal{T}(z_{kl}))\,.\end{aligned}$$

Because the orbifold action only affects the right-moving fermionic zero modes, the left-movers are untouched and Riemann's identities imply (see appendix A 2 for details)[1]

$$\sum_{\substack{a,b=0,1\\ ab=0}} (-1)^{a+b+ab}\mathcal{Z}_{a,b}\,\mathcal{W}_{a,b} = \left(\frac{\pi}{2}\right)^4 t_8F^4\,. \qquad \text{(II.27)}$$

Notice that a contribution with less than four fermionic contractions vanishes. We then obtain for the expression of the amplitude

$$\begin{aligned}\mathcal{M}^{(6)}_{(4,0)II} &= -\mathcal{N}\,\frac{1}{2}\left(\frac{\pi}{2}\right)^4 t_8F^4 \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{\frac{D-6}{2}}} \int_{\mathcal{T}} \prod_{1\le i<j\le 4} \frac{d^2\nu_i}{\tau_2}\,e^{\mathcal{Q}}\;\times \qquad \text{(II.28)}\\ &\quad\times \frac{1}{2|G|} \sum_{\substack{\bar a,\bar b=0,1\\ g,h}} (-1)^{\bar a+\bar b+\mu\bar a\bar b}(-1)^{C(\bar a,\bar b,g,h)}\,\bar{\mathcal{Z}}^{h,g}_{\bar a,\bar b}\bar{\mathcal{W}}_{\bar a,\bar b}\,.\end{aligned}$$

For the class of asymmetric $\mathbb{Z}_N$ orbifolds with n_v vector multiplets studied here, the partition function $\mathcal{Z}^{(asym)}_{a,b} = |G|^{-1}\sum_{g,h}(-1)^{C(a,b,g,h)}\mathcal{Z}^{g,h}_{a,b}$ has the following low energy expansion

$$\mathcal{Z}^{(asym)}_{0,0} = \frac{1}{\sqrt{q}} + n_v + 2 + O(q)\,; \qquad \mathcal{Z}^{(asym)}_{0,1} = \frac{1}{\sqrt{q}} - (n_v+2) + O(q)\,; \qquad \mathcal{Z}^{(asym)}_{1,0} = 0 + O(q)\,. \qquad \text{(II.29)}$$

Because the four dimensional fermionic zero modes are not saturated we have $\mathcal{Z}^{asym}_{1,1} = 0$.

Since in those constructions no massless mode arises in the twisted $h \neq 0$ sector, this sector decouples. Hence, at the $O(\bar q)$ order one has the following relation:

$$\sum_{\bar a,\bar b=0,1} (-1)^{a+b+ab}\bar{\mathcal{Z}}^{(asym)}_{\bar a,\bar b}\bar{\mathcal{W}}_{\bar a,\bar b} \to (\bar{\mathcal{W}}_{0,0} - \bar{\mathcal{W}}_{0,1})|_{\sqrt{q}} + (n_v+2)(\bar{\mathcal{W}}_{0,0} + \bar{\mathcal{W}}_{0,1})|_{q^0}\,. \qquad \text{(II.30)}$$

The massless states contribution to the field theory amplitude is given by

$$\mathcal{A}^{(n_v)}_{(4,0)II} = \frac{1}{4}\left(\frac{\pi}{2}\right)^4 t_8F^4\,\left((\bar{\mathcal{W}}_{0,0} - \bar{\mathcal{W}}_{0,1})|_{\sqrt{q}} + (n_v+2)(\bar{\mathcal{W}}_{0,0} + \bar{\mathcal{W}}_{0,1})|_{q^0}\right)\,. \qquad \text{(II.31)}$$

---

[1] $t_8F^4 = 8\mathrm{tr}(F^{(1)}F^{(2)}F^{(3)}F^{(4)}) - 2\mathrm{tr}(F^{(1)}F^{(2)})\mathrm{tr}(F^{(3)}F^{(4)}) + perms(2,3,4)$, where the traces are taken over the Lorentz indices. Setting the coupling constant to one, $t_8F^4 = stA^{tree}(1,2,3,4)$ where $A^{tree}(1,2,3,4)$ is the color stripped ordered tree amplitude between four gluons.

Using the Riemann identity (II.27) we can rewrite this expression in the following form

$$\mathcal{A}^{(n_v)}_{(4,0)II} = \frac{1}{4}\left(\frac{\pi}{2}\right)^4 t_8F^4\left(\left(\frac{\pi}{2}\right)^4 t_8\tilde{F}^4 + (n_v-6)(\bar{\mathcal{W}}_{0,0}+\bar{\mathcal{W}}_{0,1})|_{q^0} + 16\bar{\mathcal{W}}_{1,0}|_{q^0}\right) . \qquad \text{(II.32)}$$

Higher powers of $\bar{q}$ in $\mathcal{W}_{a,b}$ or in $\mathcal{Q}$ are suppressed in the field theory limit discussed in section IV.

At this level, this expression is not identical to the one derived in (II.17) from the heterotic construction. The type II and heterotic $(4,0)$ string models with n_v vector multiplets are dual to each over under the transformation $S \to -1/S$ where S is the axion-dilaton scalar in the $\mathcal{N}=4$ supergravity multiplet. We will see later that for the four-graviton amplitudes we obtain the same answer after integration.

We illustrate this analysis on the examples of the asymmetric orbifold with six or zero vector multiplets.

1. *Example: a model with six vector multiplets*

We compute now the partition function of the asymmetric orbifold obtained by the action of the right-moving fermion counting operator $(-1)^{F_R}$ and a $\mathbb{Z}_2$ action on the torus T^6 [23, 40]. The effect of the $(-1)^{F_R}$ orbifold is to project out the sixteen vector multiplets arising from the R/R sector, while preserving supersymmetry on the right moving sector. The moduli space of the theory is given by (II.1) with $n_v = 6$ and $\Gamma = \Gamma(2)$ (see [23] for instance).

The partition function for the $(4,0)$ CFT $\mathbb{Z}_2$ asymmetric orbifold model $\mathcal{Z}^{(asym),(n_v=6)}_{a,b} = \frac{1}{2}\sum_{g,h}\mathcal{Z}^{g,h}_{a,b}$ with

$$\mathcal{Z}^{h,g}_{a,b}(w) := (-1)^{ag+bh+gh}\mathcal{Z}_{a,b}\Gamma_{(4,4)}\,\Gamma^{w}_{(2,2)}\begin{bmatrix}h\\g\end{bmatrix} , \qquad \text{(II.33)}$$

where the shifted lattice sum $\Gamma^{w}_{(2,2)}\begin{bmatrix}h\\g\end{bmatrix}$ is given in [23] and recalled in appendix B. The chiral blocks $\mathcal{Z}_{a,b}$ have been defined in (II.20) and $\Gamma_{(4,4)}$ is the lattice sum of the T^4. Using the fact that $\Gamma^{w}_{(2,2)}\begin{bmatrix}h\\g\end{bmatrix}$ reduces to 0 for $h=1$, to 1 for $h=0$ and that $\Gamma_{(4,4)}\to 1$ in the field theory limit, we see that the partition function is unchanged in the sectors $(a,b)=(0,0)$ and $(0,1)$ while for the $(a,b)=(1,0)$ sector, the $(-1)^{ag}$ in (II.33) cancels the partition function when summing over g. One obtains the following result :

$$\mathcal{Z}^{(asym),(n_v=6)}_{0,0} = \mathcal{Z}_{0,0}\,; \qquad \mathcal{Z}^{(asym),(n_v=6)}_{0,1} = \mathcal{Z}_{0,1}\,; \qquad \mathcal{Z}^{(asym),(n_v=6)}_{1,0} = 0\,. \qquad \text{(II.34)}$$

Using (A.6), one checks directly that it corresponds to (II.29) with $n_v = 6$.

2. Example: models with zero vector multiplets

Now we consider the type II asymmetric orbifold models with zero vector multiplets constructed in [20] and discussed in [39].

These models are compactifications of type II superstring on a six dimensional torus with an appropriate choice for the value of the metric G_{ij} and B-field B_{ij}. The Narain lattice is given by $\Gamma^{DH} = \{p_L, p_R; p_L, p_R \in \Lambda_W(\mathfrak{g}), p_L - p_R \in \Lambda_R(\mathfrak{g})\}$ where $\Lambda_R(\mathfrak{g})$ is the root lattice of a simply laced semi-simple Lie algebra $\mathfrak{g}$, and $\Lambda_W(\mathfrak{g})$ is the weight lattice.

The asymmetric orbifold action is given by $|p_L, p_R\rangle \to e^{2i\pi p_L \cdot v_L}\, |p_L, g_R p_R\rangle$ where g_R is an element of the Weyl group of $\mathfrak{g}$ and v_L is a shift vector well chosen to avoid any massless states in the twisted sector [20, 39]. With such a choice of shift vector and because the asymmetric orbifold action leaves p_L invariant, we have $(4,0)$ supergravity models without any vector multiplets in four dimensions.

The partition function is given by

$$\mathcal{Z}^{asym}_{\bar{a},\bar{b}} = \frac{\theta\left[\substack{\bar{a}\\ \bar{b}}\right]}{\eta(\bar{\tau})^3} \frac{1}{|G|} \sum_{\{g_j, h_j\}} \prod_{i=1}^{3} \mathcal{Z}^{h_j, g_j}_{\bar{a},\bar{b}} \,, \tag{II.35}$$

where the sum runs over the sectors of the orbifold. For instance, in the $\mathbb{Z}_9$ model of Dabholkar-Harvey, one has $g_j \in j \times \{\frac{2}{9}, \frac{4}{9}, \frac{8}{9}\}$ with $j = 0, \ldots, 8$ and the same for h_j. The twisted conformal blocks are:

$$\mathcal{Z}^{h,g}_{\bar{a},\bar{b}} = \begin{cases} \left(\frac{\theta\left[\substack{\bar{a}\\ \bar{b}}\right]}{\eta(\bar{\tau})}\right)^3 \times \left(\frac{1}{\eta(\bar{\tau})}\right)^6 & \text{if } (g,h) = (0,0) \mod 2, \\ e^{i\pi a(g/2 - b/2)} 2\sin(\frac{\pi g}{2}) \frac{\theta\left[\substack{a+h\\ b+g}\right]}{\theta\left[\substack{1+h\\ 1+g}\right]} & \forall (g,h) \neq (0,0) \mod 2. \end{cases} \tag{II.36}$$

The phase in (II.19) is determined by modular invariance to be $C(\bar{a}, \bar{b}, g_R, h_R) = \sum_{i=1}^{3} (\bar{a} g^i_R + \bar{b} h^i_R + g^i_R h^i_R)$.

In the field theory limit, we perform the low energy expansion of this partition function and we find that it takes the following form for all of the models in [20, 39]:

$$\mathcal{Z}_{0,0}^{(asym),(n_v=0)} = \frac{1}{\sqrt{q}}+2+O(q)\,; \quad \mathcal{Z}_{0,1}^{(asym),(n_v=0)} = \frac{1}{\sqrt{q}}-2+O(q)\,; \quad \mathcal{Z}_{1,0}^{(asym),(n_v=0)} = 0+O(q)\,, \tag{II.37}$$

which is (II.29) with $n_v = 0$ as expected.

C. Type II Symmetric orbifold

In this section we consider $\mathcal{N} = 4$ supergravity realized as $(2,2)$ models in four dimensions. These models can be obtained from the compactification of type II string on symmetric orbifolds of $K_3 \times T^2$. The difference with the heterotic model considered in section II A is that the scalar parametrizing the coset space $SU(1,1)/U(1)$ is the Kähler modulus of the two-torus T^2 for the type IIA case or complex structure modulus for the type IIB case. The duality relation between these two models is discussed in details in [23] and [34]. These models are related to the CHL heterotic models of section II A by the non-perturbative duality relation exchanging the geometric moduli of the T^2 and the S modulus. The non-perturbative duality exchanges the heterotic T^6/G models with the type II $(K_3 \times T^2)/G$ models as given in [34, table 2].

Models with $n_v \geq 2$ have been constructed in [31]. The model with $n_v = 22$ is an orbifold $T^4/\mathbb{Z}_2 \times T^2$, and the following models with $n_v \in \{14, 10\}$ are successive $\mathbb{Z}_2$ orbifolds of the first one. The model with $n_v = 6$ is a freely acting $\mathbb{Z}_2$ orbifold of the $T^4/\mathbb{Z}_2 \times T^2$ theory that simply projects out the sixteen vector multiplets of the R/R sector. The four-graviton amplitude can be effectively written in terms of the (g,h) sectors of the first $\mathbb{Z}_2$ orbifold of the T^4, and writes

$$\begin{aligned}\mathcal{M}_{(2,2)}^{(n_v)} = \mathcal{N} \int_{\mathcal{F}} \frac{d^2\tau}{\tau_2^{\frac{D-6}{2}}} \int_{\mathcal{T}} \prod_{1\leq i<j\leq 4} \frac{d^2\nu_i}{\tau_2}\, e^{\mathcal{Q}} \; \times \qquad\qquad (\text{II.38})\\ \times \frac{1}{4|G|} \sum_{h,g=0}^{1} \sum_{\substack{a,b=0,1\\ \bar a,\bar b=0,1}} (-1)^{a+b+ab}(-1)^{\bar a+\bar b+\bar a\bar b}\, \mathcal{Z}_{a,b}^{h,g,(n_v)}\, \bar{\mathcal{Z}}_{\bar a,\bar b}^{h,g,(n_v)}\, \mathcal{W}_{a,b}\bar{\mathcal{W}}_{\bar a,\bar b}\,,\end{aligned}$$

where $\mathcal{N}$ is the same overall normalization as for the previous amplitudes and $\mathcal{Z}_{a,b}^{h,g,(n_v)}$ is defined in appendix B. Forgetting about the lattice sum that anyway reduces to 1 in the field theory limit,

$$\mathcal{Z}^{h,g}_{a,b} = c_h \frac{\theta\left[{a \atop b}\right]^2(0|\tau)\,\theta\left[{a+h \atop b+g}\right](0|\tau)\,\theta\left[{a-h \atop b-g}\right](0|\tau)}{\eta(\tau)^6\,\theta\left[{1+h \atop 1+g}\right]^2} = c_h(-1)^{(a+h)g}\left(\frac{\theta\left[{a \atop b}\right](0|\tau)\,\theta\left[{a+h \atop b+g}\right](0|\tau)}{\eta(\tau)^3\,\theta\left[{1+h \atop 1+g}\right]}\right)^2, \tag{II.39}$$

where c_h is an effective number whose value depend on h in the following way: $c_0 = 1$ and $c_1 = \sqrt{n_v - 6}$. This number represents the successive halving of the number of twisted R/R states. We refer to appendix B for details.

The sum over the spin structure in the untwisted sector $(g,h) = (0,0)$ is performed as before using Riemann's identities:

$$\sum_{\substack{a,b=0,1\\ ab=0}} (-1)^{a+b+ab}\mathcal{Z}^{0,0}_{a,b}\mathcal{W}_{a,b} = \left(\frac{\pi}{2}\right)^4 t_8F^4\,. \tag{II.40}$$

In the twisted sectors $(h,g) \neq (0,0)$ we remark that $\mathcal{Z}^{0,1}_{0,1} = Z^{0,1}_{0,0}$, $\mathcal{Z}^{1,0}_{1,0} = \mathcal{Z}^{1,0}_{0,0}$, $\mathcal{Z}^{1,1}_{1,0} = \mathcal{Z}^{1,1}_{0,1}$, and $\mathcal{Z}^{1,0}_{0,1} = \mathcal{Z}^{0,1}_{1,0} = \mathcal{Z}^{1,1}_{0,0} = 0$, which gives for the chiral blocks in (II.38):

$$\begin{aligned}
\sum_{\substack{a,b=0,1\\ ab=0}} (-1)^{a+b+ab}\mathcal{Z}^{0,1}_{a,b}\mathcal{W}_{a,b} &= \mathcal{Z}^{0,1}_{0,0}\,\left(\mathcal{W}_{0,0} - \mathcal{W}_{0,1}\right),\\
\sum_{\substack{a,b=0,1\\ ab=0}} (-1)^{a+b+ab}\mathcal{Z}^{1,0}_{a,b}\mathcal{W}_{a,b} &= \mathcal{Z}^{1,0}_{0,0}\,\left(\mathcal{W}_{0,0} - \mathcal{W}_{1,0}\right),\\
\sum_{\substack{a,b=0,1\\ ab=0}} (-1)^{a+b+ab}\mathcal{Z}^{1,1}_{a,b}\mathcal{W}_{a,b} &= \mathcal{Z}^{1,1}_{0,1}\,\left(\mathcal{W}_{0,1} - \mathcal{W}_{1,0}\right)\,.
\end{aligned} \tag{II.41}$$

Therefore the contributions in the correlator take the simplified form

$$\begin{aligned}
&\frac{1}{4|G|}\sum_{g,h}\sum_{\substack{a,b=0,1\\ \bar a,\bar b=0,1}} (-1)^{a+b+ab}(-1)^{\bar a+\bar b+\bar a\bar b}\mathcal{Z}^{h,g}_{a,b}\bar{\mathcal{Z}}^{\bar h,\bar g}_{\bar a,\bar b}\,\mathcal{W}_{a,b}\bar{\mathcal{W}}_{\bar a,\bar b}\\
&= \frac{1}{8}\left(\frac{\pi}{2}\right)^8 t_8t_8R^4 + \frac{1}{8}\,\left|\mathcal{Z}^{0,1}_{0,0}(\mathcal{W}_{0,0} - \mathcal{W}_{0,1})\right|^2 + \frac{1}{8}\,|\mathcal{Z}^{1,0}_{0,0}(\mathcal{W}_{0,0} - \mathcal{W}_{1,0})|^2 + \frac{1}{8}\,|\mathcal{Z}^{1,1}_{0,1}(\mathcal{W}_{0,1} - \mathcal{W}_{1,0})|^2\,,
\end{aligned} \tag{II.42}$$

where $t_8t_8R^4$ is the Lorentz scalar built from four powers of the Riemann tensor arising at the linearized level as the product $t_8t_8R^4 = t_8F^4\,t_8\tilde F^4$.[2]

Since the conformal blocks $\mathcal{Z}^{h,g}_{a,b}$ have the q-expansion (see (A.3))

$$\mathcal{Z}^{0,1}_{0,0} = \frac{1}{\sqrt{q}} + 4\sqrt{q} + O(q)\,;\ \mathcal{Z}^{1,0}_{0,0} = 4\sqrt{n_v - 6} + O(q)\,;\ \mathcal{Z}^{1,1}_{0,1} = 4\sqrt{n_v - 6} + O(q)\,, \tag{II.43}$$

---

[2] This Lorentz scalar is the one obtained from the four-graviton tree amplitude $t_8t_8R^4 = stuM^{tree}(1,2,3,4)$ setting Newton's constant to one.

the massless contribution to the integrand of (II.42) is given by

$$\mathcal{A}^{(n_v)}_{(2,2)} = \frac{1}{8}\left(\frac{\pi}{2}\right)^8 t_8 t_8 R^4 + \frac{1}{8}\left|\mathcal{W}_{0,0}|_{\sqrt{q}} - \mathcal{W}_{0,1}|_{\sqrt{q}}\right|^2 \\ + 2(n_v - 6)\left(|\mathcal{W}_{0,0}|_{q^0} - \mathcal{W}_{1,0}|_{q^0}|^2 + |\mathcal{W}_{0,1}|_{q^0} - \mathcal{W}_{1,0}|_{q^0}|^2\right) . \quad \text{(II.44)}$$

We notice that the bosonic piece $\mathcal{W}^B$ in $\mathcal{W}_{a,b}$ in (II.24) cancels in each term of the previous expression, due to the minus sign between the $\mathcal{W}_{a,b}$'s in the squares.

The integrand of the four-graviton amplitude takes a different form in the $(2,2)$ construction compared to the expression for the $(4,0)$ constructions in heterotic in (II.17) and asymmetric type II models in (II.32). We will show that even after taking the limit and performing the integrals the amplitudes will be different.[3]

This is not really surprising since for a given number vector multiplets n_v the type II $(2,2)$ models are U-dual to the $(4,0)$ models, and again this duality is non-perturbative. The relations between the various string theory models are discussed in the next section.

III. COMPARISON OF THE STRING MODELS

A. Massless spectrum

The spectrum of type II superstring in ten dimensions is given by the following GSO sectors: the graviton G_{MN}, the B-field B_{MN}, and the dilaton Φ come from the NS/NS sector, the gravitini ψ^M, and the dilatini λ come from the R/NS and NS/R sectors, while the one-form C_M and three-form C_{MNP} come from the R/R sector in type the IIA string. The dimensional reduction of type II string on a torus preserves all of the thirty-two supercharges, and leads to the $\mathcal{N} = 8$ supergravity multiplet in four dimensions.

The reduction to $\mathcal{N} = 4$ supersymmetry (preserving sixteen supercharges) leads to the following content. The NS/NS sector contributes to the $\mathcal{N} = 4$ supergravity multiplet and six vector supermultiplets. The R/R sector contributes to the $\mathcal{N} = 4$ spin $\frac{3}{2}$ multiplet and vector supermultiplets with a multiplicity depending on the model.

The first Riemann vanishing identity

$$\sum_{a,b=0,1} (-1)^{a+b+ab} \mathcal{Z}_{a,b} = 0 \,, \quad \text{(III.1)}$$

---

[3] In the previous analysis of $(4,0)$ models, (II.17) and (II.32) were also different, but they are equal in the field theory limit as was already said.

reflects the action of the $\mathcal{N}=4$ supersymmetry inside the one-loop amplitudes.

The q-expansion of this identity gives

$$\left(\frac{1}{\sqrt{q}}+8+O(\sqrt{q})\right)-\left(\frac{1}{\sqrt{q}}-8+O(\sqrt{q})\right)-(16+O(q))=0\,. \tag{III.2}$$

The first two terms are the expansion of $\mathcal{Z}_{0,0}$ and $\mathcal{Z}_{0,1}$, the last one is the expansion of $\mathcal{Z}_{1,0}$. The cancellation of the $1/\sqrt{q}$ terms shows that the GSO projection eliminates the tachyon from the spectrum, and at the order q^0 the cancellation results in the matching between the bosonic and fermionic degrees of freedom.

In amplitudes, chiral $\mathcal{N}=4$ supersymmetry implies the famous Riemann identities, stating that $0 \leq n \leq 3$ points one-loop amplitude vanish (see eq. (A.25)), while at four points we have

$$\sum_{a,b=0,1}(-1)^{a+b+ab}\mathcal{Z}_{a,b}\mathcal{W}_{a,b}=\left(\frac{\pi}{2}\right)^4 t_8F^4\,. \tag{III.3}$$

The spin independent part of $\mathcal{W}^B$ cancels trivially due to (III.1) as well as the contributions with less than four fermionic contractions $\mathcal{S}_{2;a,b}$ in $\mathcal{W}^F_{a,b}$ in (II.25).

The cancellation of the tachyon yields at the first order in the q-expansion of (III.3)

$$\mathcal{W}_{0,0}|_{q^0}-\mathcal{W}_{0,1}|_{q^0}=0\,. \tag{III.4}$$

The next term in the expansion gives an identity describing the propagation of the super-Yang-Mills multiplet in the loop

$$8(\mathcal{W}_{0,0}|_{q^0}+\mathcal{W}_{0,1}|_{q^0}-2\mathcal{W}_{1,0}|_{q^0})+(\mathcal{W}_{0,0}|_{\sqrt{q}}-\mathcal{W}_{0,1}|_{\sqrt{q}})=\left(\frac{\pi}{2}\right)^4 t_8F^4\,. \tag{III.5}$$

In this equation, one should have vectors, spinors and scalars propagating according to the sector of the theory. In $\mathcal{W}_{a,b}$, $a=0$ is the NS sector, and $a=1$ is the Ramond sector. The scalars have already been identified in (III.4) and correspond to $\mathcal{W}_{0,0}|_{q^0}+\mathcal{W}_{0,1}|_{q^0}$. The vector, being a massless bosonic degree of freedom should then correspond to $\mathcal{W}_{0,0}|_{\sqrt{q}}-\mathcal{W}_{0,1}|_{\sqrt{q}}$. Finally, the fermions correspond to $\mathcal{W}_{1,0}|_{q^0}$. The factor 8 in front of the first term is the number of degree of freedom of a vector in ten dimension, one can check that the number of bosonic degrees of freedom match the number of fermionic degrees of freedom.

B. Amplitudes and supersymmetry

In this section we discuss the relations between the four-graviton amplitudes in the various $\mathcal{N}=4$ supergravity models in the field theory limit. We apply the logic of the previous section about the spectrum of left or right movers to the tensor product spectrum and see that we can precisely identify the contributions to the amplitude, both in the $(4,0)$ and $(2,2)$ models. However, we would like to emphasize that the differences between the two models will not be visible from this spectrum point of view but will be seen in the amplitude computations of sections III C 2 and IV B.

The complete evaluation of the amplitudes will be performed in section IV.

As was said, the field theory limit is obtained by considering the large τ_2 region, and the integrand of the field theory amplitude is given by

$$A_X^{(n_v)} = \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1 \, \mathcal{A}_X^{(n_v)} \,, \tag{III.6}$$

where $X \in \{(4,0)het,\ (4,0)II,\ (2,2)\}$ depending on the model, as in (II.17), (II.32) or (II.44) respectively.

At one-loop this quantity is the sum of the contribution from n_v $\mathcal{N}=4$ vector (spin 1) supermultiplets running in the loop and the $\mathcal{N}=4$ spin-2 supermultiplet

$$A_X^{(n_v)} = A_X^{spin\ 2} + n_v\, A_X^{spin\ 1} \,. \tag{III.7}$$

For the case of the type II asymmetric orbifold models with n_v vector multiplet we deduce from (II.32)

$$A_{(4,0)II}^{spin\ 1} = \frac{1}{4}\left(\frac{\pi}{2}\right)^4 t_8F^4 \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1 \, (\mathcal{W}_{0,0}|_{q^0} + \mathcal{W}_{0,1}|_{q^0}) \,. \tag{III.8}$$

Since t_8F^4 is the supersymetric left-moving sector contribution (recall the supersymmetry identity in (III.3)), it corresponds to an $\mathcal{N}=4$ vector multiplet and we recognize in (III.8) the product of this multiplet with the scalar from the right-moving sector:

$$(1_{\mathbf{1}}, 1/2_{\mathbf{4}}, 0_{\mathbf{6}})_{\mathcal{N}=4} = (1_{\mathbf{1}}, 1/2_{\mathbf{4}}, 0_{\mathbf{6}})_{\mathcal{N}=4} \otimes (0_{\mathbf{1}})_{\mathcal{N}=0} \,. \tag{III.9}$$

This agrees with the identification made in the previous subsection where $\mathcal{W}_{0,0}|_{q^0} + \mathcal{W}_{0,1}|_{q^0}$ was argued to be a scalar contribution.

The contribution from the $\mathcal{N}=4$ supergravity multiplet running in the loop is given by

$$A_{(4,0)II}^{spin\ 2} = \frac{1}{4}\left(\frac{\pi}{2}\right)^4 t_8F^4 \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1 \left[2(\mathcal{W}_{0,0}|_{q^0} + \mathcal{W}_{0,1}|_{q^0}) + (\mathcal{W}_{0,0}|_{\sqrt{q}} - \mathcal{W}_{0,1}|_{\sqrt{q}})\right] . \tag{III.10}$$

The factor of 2 is number of degrees of freedom of a vector in four dimensions. Since $\mathcal{Z}_{1,0}^{asym} = 0 + O(q)$ for the $(4,0)$ model asymmetric orbifold construction, the integrand of the four gravitons amplitude in (II.29) does not receive any contribution from the right moving R sector. Stated differently, the absence of $\mathcal{W}_{1,0}$ implies that both R/R and NS/R sectors are projected out, leaving only the contribution from the NS/NS and R/NS. Thus, four $\mathcal{N}=4$ spin $\frac{3}{2}$ supermultiplet and sixteen $\mathcal{N}=4$ spin 1 supermultiplet are projected out, leaving at most six vector multiplets. This number is decreased to zero in the Dabholkar-Harvey construction [20].

From (III.10) we recognize that the $\mathcal{N}=4$ supergravity multiplet is obtained by the following tensor product

$$(2_{\mathbf{1}}, 3/2_{\mathbf{4}}, 1_{\mathbf{6}}, 1/2_{\mathbf{4}}, 0_{\mathbf{2}})_{\mathcal{N}=4} = (1_{\mathbf{1}}, 1/2_{\mathbf{4}}, 0_{\mathbf{6}})_{\mathcal{N}=4} \otimes (1_{\mathbf{1}})_{\mathcal{N}=0}\,. \tag{III.11}$$

The two real scalars arise from the trace part and the anti-symmetric part (after dualisation in four dimensions) of the tensorial product of the two vectors. Using the identification of $\mathcal{W}_{0,0}|_{q^0} + \mathcal{W}_{0,1}|_{q^0}$ with a scalar contribution and the equation (A.30) we can now identify $\mathcal{W}_{0,0}|_{\sqrt{q}} - \mathcal{W}_{0,0}|_{\sqrt{q}}$ with the contribution of a vector and two scalars. This confirms the identification of $\mathcal{W}_{1,0}|_{q^0}$ with a spin $\frac{1}{2}$ contribution in the end of section III A.

Since

$$(3/2_{\mathbf{1}}, 1_{\mathbf{4}}, 1/2_{\mathbf{6}+\mathbf{1}}, 0_{\mathbf{4}+\bar{\mathbf{4}}})_{\mathcal{N}=4} = (1_{\mathbf{1}}, 1/2_{\mathbf{4}}, 0_{\mathbf{6}})_{\mathcal{N}=4} \otimes (1/2)_{\mathcal{N}=0}\,, \tag{III.12}$$

we see that removing the four spin $\frac{1}{2}$ (that is, the $\mathcal{W}_{1,0}|_{q^0}$ contribution) of the right-moving massless spectrum of the string theory construction in asymmetric type II models removes the contribution from the massless spin $\frac{3}{2}$ to the amplitudes. For the asymmetric type II model, using (III.5), we can present the contribution from the $\mathcal{N}=4$ supergravity multiplet in a form that reflects the decomposition in the $\mathcal{N}=8$ supergravity multiplet into $\mathcal{N}=4$ supermultiplets

$$\begin{aligned}(2_{\mathbf{1}}, 3/2_{\mathbf{8}}, 1_{\mathbf{28}}, 1/2_{\mathbf{56}}, 0_{\mathbf{70}})_{\mathcal{N}=8} &= (2_{\mathbf{1}}, 3/2_{\mathbf{4}}, 1_{\mathbf{6}}, 1/2_{\mathbf{4}}, 0_{\mathbf{1}+\mathbf{1}})_{\mathcal{N}=4}\\ &\oplus\ \mathbf{4}\,(3/2_{\mathbf{1}}, 1_{\mathbf{4}}, 1/2_{\mathbf{6}+\mathbf{1}}, 0_{\mathbf{4}+\bar{\mathbf{4}}})_{\mathcal{N}=4}\\ &\oplus\ \mathbf{6}\,(1_{\mathbf{1}}, 1/2_{\mathbf{4}}, 0_{\mathbf{6}})_{\mathcal{N}=4}\,,\end{aligned} \tag{III.13}$$

as

$$A^{spin\ 2}_{(4,0)II} = A^{spin\ 2}_{\mathcal{N}=8} - 6\,A^{spin\ 1}_{(4,0)II} - 4\,A^{spin\ \frac{3}{2}}_{(4,0)II}\,, \tag{III.14}$$

where we have introduced the $\mathcal{N}=8$ spin 2 supergravity contribution

$$A^{spin\ 2}_{\mathcal{N}=8} = \frac{1}{4}\left(\frac{\pi}{2}\right)^{8} t_8 t_8 R^4\,, \tag{III.15}$$

and the $\mathcal{N}=4$ spin $\frac{3}{2}$ supergravity contribution

$$A^{spin\ \frac{3}{2}}_{(4,0)II} = -\left(\frac{\pi}{2}\right)^4 t_8F^4 \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, \mathcal{W}_{1,0}|_{q^0}\,. \tag{III.16}$$

For the $(2,2)$ models the massless states contributions to the amplitude are given in (II.44). The contribution from the vector multiplet is

$$A^{spin\ 1}_{(2,2)} = 2\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, \left(\left|\mathcal{W}_{0,0}|_{q^0} - \mathcal{W}_{1,0}|_{q^0}\right|^2 + \left|\mathcal{W}_{0,1}|_{q^0} - \mathcal{W}_{1,0}|_{q^0}\right|^2\right)\,. \tag{III.17}$$

Using that $\left|\mathcal{W}_{0,0}|_{q^0} - \mathcal{W}_{0,1}|_{q^0}\right|^2 = \left|\mathcal{W}_{1,0}|_{q^0} - \mathcal{W}_{0,1}|_{q^0}\right|^2 = \frac{1}{4}\left|\mathcal{W}_{0,0}|_{q^0} + \mathcal{W}_{0,1}|_{q^0} - 2\mathcal{W}_{1,0}|_{q^0}\right|^2$ as a consequence of (III.4), we can rewrite this as

$$A^{spin\ 1}_{(2,2)} = \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{1\leq i<j\leq 4} d\nu_i^1\, \left(\left|\mathcal{W}_{0,0}|_{q^0} + \mathcal{W}_{1,0}|_{q^0} - 2\mathcal{W}_{0,1}|_{q^0}\right|^2\right)\,, \tag{III.18}$$

showing that this spin 1 contribution in the $(2,2)$ models arises as the product of two $\mathcal{N}=2$ hypermultiplets $Q=(2\times 1/2_{\mathbf{1}}, 2\times 0_{\mathbf{2}})_{\mathcal{N}=2}$

$$2\times(1_{\mathbf{1}}, 1/2_{\mathbf{4}}, 0_{\mathbf{6}})_{\mathcal{N}=4} = (2\times 1/2_{\mathbf{1}}, 2\times 0_{\mathbf{2}})_{\mathcal{N}=2} \otimes (2\times 1/2_{\mathbf{1}}, 2\times 0_{\bar{\mathbf{2}}})_{\mathcal{N}=2}\,. \tag{III.19}$$

The contribution from the $\mathcal{N}=4$ supergravity multiplet running in the loop (obtained from (II.44) by setting $n_v=0$) can be presented in a form reflecting the decomposition in (III.13)

$$A^{spin\ 2}_{(2,2)} = A^{spin\ 2}_{\mathcal{N}=8} - 6A^{spin\ 1}_{(2,2)} - 4\,A^{spin\ \frac{3}{2}}_{(2,2)}\,, \tag{III.20}$$

where $A^{spin\ \frac{3}{2}}_{(2,2)}$ is given by

$$A^{spin\ \frac{3}{2}}_{(2,2)} = -\frac{1}{8}\,A^{spin\ 2}_{\mathcal{N}=8} - \frac{1}{32}\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{1\leq i<j\leq 4} d\nu_i^1\, \left|\mathcal{W}_{0,0}|_{\sqrt{q}} - \mathcal{W}_{0,1}|_{\sqrt{q}}\right|^2. \tag{III.21}$$

C. Comparing of the string models

The integrands of the amplitudes in the two $(4,0)$ models in (II.17) and (II.32) or in the $(2,2)$ models in (II.44) take a different form. In this section we show first the equality between the integrands of the $(4,0)$ models, then that the difference with the $(2,2)$ models can be attributed to the contribution of the vector multiplets.

The comparison is done in the field theory limit where $\tau_2\to+\infty$ and $\alpha'\to 0$ with $t=\alpha'\tau_2$ hold fixed. In the field theory limit the real parts of the ν_i variables are integrated over the range $-\frac{1}{2} \leq \nu_i^1 \leq \frac{1}{2}$. In this limit the position of the vertex operators scale as $\nu_i = \nu_i^1 + i\tau_2\omega_i$. The positions of the external legs on the loop are then denoted by $0 \leq \omega_i \leq 1$ and are ordered according to the kinematical region under consideration. In this section we discuss the real part integration, the integration over the ω_i variables will be performed in section IV.

1. *Comparing the* (4,0) *models*

In the heterotic string amplitude (II.17), we can identify two distinct contributions; n_v vector multiplets and one $\mathcal{N} = 4$ supergravity multiplet running in the loop. The contribution of the vector multiplets writes

$$A^{spin\ 1}_{(4,0)het} = \frac{1}{2}\left(\frac{\pi}{2}\right)^4 t_8F^4 \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, \bar{\mathcal{W}}^B|_{\bar{q}^0}\,, \qquad \text{(III.22)}$$

and for the supergravity multiplet we have at the leading order in α'

$$A^{spin\ 2}_{(4,0)het} = \frac{1}{2}\left(\frac{\pi}{2}\right)^4 t_8F^4 \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1 \left((\bar{\mathcal{W}}^B|_{\bar{q}}(1+\alpha'\delta Q) + \bar{\mathcal{W}}^B|_{\bar{q}^0}\mathcal{Q}|_{\bar{q}} + 2\bar{\mathcal{W}}^B|_{\bar{q}^0}\right). \qquad \text{(III.23)}$$

The vector multiplet contributions take different forms in the heterotic construction in (III.22) and the type II models in (III.8). However using the expansion of the fermionic propagators given in appendix A 2, it is not difficult to perform the integration over ν_i^1 in (III.8). We see that

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{1\leq i<j\leq 4} d\nu_i^1\, (\mathcal{W}^F_{0,0}|_{\bar{q}^0} + \mathcal{W}^F_{0,1}|_{\bar{q}^0}) = 0\,. \qquad \text{(III.24)}$$

Thus there only remain the bosonic part of $\mathcal{W}_{a,b}$, and we find that the contribution of the vector multiplet is the same in the heterotic and asymmetric orbifold constructions

$$A^{spin\ 1}_{(4,0)het} = A^{spin\ 1}_{(4,0)II}\,. \qquad \text{(III.25)}$$

The case of the $\mathcal{N} = 4$ super-graviton is a little more involved. In order to simplify the argument we make a choice of helicity that makes the expressions more manageable.

We make the following choice of helicity $(1^{++}, 2^{++}, 3^{--}, 4^{--})$ and reference momenta $q_1 = q_2 = k_3$ and $q_3 = q_4 = k_1$. In that manner the covariant quantities t_8F^4 and $t_8t_8R^4$ write respectively in the spinor helicity formalism[4] $2t_8F^4 = \langle k_1\, k_2\rangle^2\, [k_3\, k_4]^2$, and $4t_8t_8R^4 =$

---

[4] A null vector $k^2 = 0$ is parametrized as $k_{\alpha\dot\alpha} = k_\alpha \bar{k}_{\dot\alpha}$ where $\alpha, \dot\alpha = 1,2$ are $SL(2,\mathbb{C})$ two dimensional spinor indices. The positive and negative helicity polarization vectors are given by $\epsilon^+(k,q)_{\alpha\dot\alpha} := \frac{q_\alpha \bar{k}_{\dot\alpha}}{\sqrt{2}\,\langle q\, k\rangle}$ and $\epsilon^-(k,q)_{\alpha\dot\alpha} := -\frac{k_\alpha \bar{q}_{\dot\alpha}}{\sqrt{2}\,[q\, k]}$ where q is a massless reference momentum. The self-dual and anti-self-dual field strengths with $F^-_{\alpha\beta} := \sigma^{mn}_{\alpha\beta}F_{mn} = \frac{k_\alpha k_\beta}{\sqrt{2}}$ and $F^+_{\dot\alpha\dot\beta} := \bar\sigma^{mn}_{\dot\alpha\dot\beta}F_{mn} = -\frac{\bar{k}_{\dot\alpha}\bar{k}_{\dot\beta}}{\sqrt{2}}$.

$\langle k_1\, k_2\rangle^4\, [k_3\, k_4]^4$. We can now simplify the various kinematical factors, $\mathcal{W}^B$ for the heterotic string and the $\mathcal{W}_{a,b}$'s for the type II models. We find $\bar{\mathcal{W}}^B = \frac{1}{2} t_8 \tilde{F}^4\, \tilde{\mathcal{W}}^B$ where

$$\tilde{\mathcal{W}}^B = \tilde{\mathcal{W}}^B_1 + \frac{1}{\alpha' u} \tilde{\mathcal{W}}^B_2\,, \tag{III.26}$$

with

$$\begin{aligned}
\tilde{\mathcal{W}}^B_1 &= (\bar\partial\mathcal{P}(12) - \bar\partial\mathcal{P}(14))(\bar\partial\mathcal{P}(21) - \bar\partial\mathcal{P}(24))(\bar\partial\mathcal{P}(32) - \bar\partial\mathcal{P}(34))(\bar\partial\mathcal{P}(42) - \bar\partial\mathcal{P}(43))\,,\\
\tilde{\mathcal{W}}^B_2 &= \bar\partial^2\mathcal{P}(24)(\bar\partial\mathcal{P}(12) - \bar\partial\mathcal{P}(14))(\bar\partial\mathcal{P}(32) - \bar\partial\mathcal{P}(34))\,.
\end{aligned} \tag{III.27}$$

In these equations it is being understood that $\mathcal{P}(ij)$ stands for $\mathcal{P}(\nu_i - \nu_j)$. We find as well that $\mathcal{W}^F_{a,b} = \frac{1}{2} t_8 F^4\, \tilde{\mathcal{W}}^F_{a,b}$ with $\tilde{\mathcal{W}}_{a,b} = \tilde{\mathcal{S}}_{4;a,b} + \tilde{\mathcal{S}}_{2;a,b}$ where

$$\begin{aligned}
\tilde{\mathcal{S}}_{4;a,b} &= \frac{1}{2^8}\left(S_{a,b}(12)^2 S_{a,b}(34)^2 - S_{a,b}(1234) - S_{a,b}(1432) - S_{a,b}(1423)\right)\\
\tilde{\mathcal{S}}_{2;a,b} &= \frac{1}{2^4}(\partial\mathcal{P}(12) - \partial\mathcal{P}(14))(\partial\mathcal{P}(21) - \partial\mathcal{P}(24)) S_{a,b}(34)^2\\
&\quad + \frac{1}{2^4}(\partial\mathcal{P}(32) - \partial\mathcal{P}(34))(\partial\mathcal{P}(42) - \partial\mathcal{P}(43)) S_{a,b}(12)^2\,,
\end{aligned} \tag{III.28}$$

where we have used a shorthand notation; $S_{a,b}(ij)$ stands for $S_{a,b}(z_i - z_j)$ while $S_{a,b}(ijkl)$ stands for $S_{a,b}(z_i - z_j) S_{a,b}(z_j - z_k) S_{a,b}(z_k - z_l) S_{a,b}(z_l - z_i)$.

Let us come back to the comparison of the $\mathcal{N} = 4$ spin 2 multiplet contributions in both the type II asymmetric orbifold model in (III.10) and the heterotic construction in (III.23).

We consider the following part of (III.23)

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu^1_i\, (\bar{\mathcal{W}}^B|_{\bar q}(1 + \alpha'\delta Q) + \bar{\mathcal{W}}^B|_{\bar q^0}\, \mathcal{Q}|_{\bar q})\,, \tag{III.29}$$

defined in the field theory limit for large τ_2.

The integral over the ν^1_i will kill any term that have a non zero phase $e^{i\pi(a\nu^1_1 + b\nu^1_2 + c\nu^1_3 + d\nu^1_4)}$ where a, b, c, d are non all-vanishing integers. From $\tilde{W}^B_1$ we have terms of the form $\partial\delta P(ij) \times \partial\mathcal{P}(ji)|_{\bar q} \times (\partial P(kl) - \partial P(k'l'))(\partial P(rs) - \partial P(r's'))$. Using the definition for $\delta\mathcal{P}(ij)$ given in (A.15) and the order $\bar q$ coefficient of the propagator in (A.14), we find that

$$\partial\delta P(ij) \times \partial\mathcal{P}(ji)|_{\bar q} = -\frac{i\pi^2}{2} \sin(2\pi\nu_{ij})\, \mathrm{sign}(\omega_{ij}) e^{2i\pi \mathrm{sign}(\omega_{ij})\nu_{ij}}\,, \tag{III.30}$$

which integrates to $-\pi^2/4$. All such terms with $(ij) = (12)$ and $(ij) = (34)$ contribute in total to

$$\frac{1}{2}\left(\frac{\pi}{2}\right)^4 [(\partial P(12) - \partial P(14))(\partial P(21) - \partial P(24)) + (\partial P(32) - \partial P(34))(\partial P(42) - \partial P(43))]\,, \tag{III.31}$$

where here $\partial P(ij)$ is for the derivative of the propagator in the field theory $\partial P(\omega_i - \omega_j)$ given by

$$\partial P(\omega) = 2\omega - \mathrm{sign}(\omega)\,. \tag{III.32}$$

The last contribution from $\tilde{W}_1^B$ is for $(ij) = (24)$ leading to the same kind of contribution, that will actually be cancelled by terms coming from similar contractions in $\tilde{\mathcal{W}}_2^B|_{\bar q}$. More precisely, the non zero contractions involved yield

$$(\partial^2\mathcal{P}(24)e^{\mathcal{Q}})|_{\bar q} = -\frac{\alpha'\pi^2}{2}\left(\cos(2\pi\nu_{24})e^{2i\pi\mathrm{sign}(\omega_{24})\nu_{24}} - 2e^{2i\pi\mathrm{sign}(\omega_{24})\nu_{24}}\sin^2(\pi\nu_{24})\right)\,, \tag{III.33}$$

which integrates to $-\alpha'\pi^2/2$. The α' compensates the $1/\alpha'$ factor in (III.26) and this contribution precisely cancels the one from (III.30) with $(ij) = (24)$. Other types of terms with more phase factors from the propagator turn out to vanish after summation. In all, we get $-\pi^4 W_3/4$, where

$$W_3 = -\frac{1}{8}\Big((\partial P(12) - \partial P(14))(\partial P(21) - \partial P(24)) + (\partial P(32) - \partial P(34))(\partial P(42) - \partial P(43))\Big)\,, \tag{III.34}$$

Finally, let's look at the totally contracted terms of the form $\partial\delta P(ik)\partial\delta P(kl)\partial\delta P(lj) \times \partial\mathcal{P}(ij)|_{\bar q}$ that come from $\tilde{\mathcal{W}}_1^B|\bar q$. Those are the only terms of that type that survive the ν^1 integrations since they form closed chains in their arguments. They give the following terms

$$i\frac{\pi^4}{8}\sin(\pi\nu_{ij})\mathrm{sign}(\omega_{ik})\mathrm{sign}(\omega_{kl})\mathrm{sign}(\omega_{lj})e^{2i\pi\left(\mathrm{sign}(\omega_{ik})\nu_{ik}+\mathrm{sign}(\omega_{kl})\nu_{kl}+\mathrm{sign}(\omega_{lj})\nu_{lj}\right)}\,. \tag{III.35}$$

They integrate to $\pi^4/16$ if the vertex operators are ordered according to $0 \leq \omega_i < \omega_k < \omega_l < \omega_j \leq 1$ or in the reversed ordering. Hence, from $\tilde{\mathcal{W}}_1^B$ we will get one of the ordering we want in our polarization choice, namely the region (s,t). From $\tilde{\mathcal{W}}_2 e^{\mathcal{Q}}$, a similar computation gives the two other kinematical regions (s,u) and (t,u), in all we have a total integrated contribution of $\pi^4/16$. We collect all the different contributions that we have obtained, and (III.23) writes:

$$A^{spin\ 2}_{(4,0)het} = \frac{1}{4}\left(\frac{\pi}{2}\right)^8 t_8t_8R^4\left(1 - 4W_3 + 2\,W^B\right), \tag{III.36}$$

where we used that $t_8t_8R^4 = t_8F^4t_8\tilde{F}^4$ and defined

$$W^B := \left(\frac{2}{\pi}\right)^4\int_{-\frac{1}{2}}^{\frac{1}{2}}\prod_{i=1}^{4} d\nu_i^1\,\tilde{\mathcal{W}}^B|_{\bar q^0}\,, \tag{III.37}$$

evaluated in the field theory limit $\tau_2 \to \infty$. In this limit this quantity is given by $W^B = W_1 + W_2$ with

$$\begin{aligned} W_1 &= \frac{1}{16}(\partial P(12) - \partial P(14))(\partial P(21) - \partial P(24))(\partial P(32) - \partial P(34))(\partial P(42) - \partial P(43))\,, \\ W_2 &= \frac{1}{4\pi}\frac{1}{\alpha'\tau_2 u}\,\partial^2 P(24)(\partial P(12) - \partial P(14))(\partial P(32) - \partial P(34))\,, \end{aligned} \tag{III.38}$$

where $\partial^n P(\omega)$ is the n-th derivative of the field theory propagator (III.32) and where $\alpha'\tau_2$ is the proper time of the field one-loop amplitude.

We now turn to the spin 2 contribution in the type II asymmetric orbifold models given in (III.10). Using the q-expansion given in appendix A 2 c, we find that

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, (\tilde{\mathcal{W}}_{0,0}|_{\sqrt{q}} - \tilde{\mathcal{W}}_{0,1}|_{\sqrt{q}}) = \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, (\tilde{\mathcal{W}}^F_{0,0}|_{\sqrt{q}} - \tilde{\mathcal{W}}^F_{0,1}|_{\sqrt{q}}) = 2\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, \tilde{\mathcal{W}}^F_{0,0}|_{\sqrt{q}}\,. \tag{III.39}$$

We have then terms of the form $\tilde{\mathcal{S}}_{2;0,0}$ and $\tilde{\mathcal{S}}_{4;0,0}$. Their structure is similar to the terms in the heterotic case with respectively two and four bosonic propagators contracted. The bosonic propagators do not have a $\sqrt{q}$ piece and since $\tilde{\mathcal{S}}_{0,0}(12)^2|_{\sqrt{q}} = \tilde{\mathcal{S}}_{0,0}(34)^2|_{\sqrt{q}} = 4\pi^2$ we find that the terms in $\mathcal{S}_{2;0,0}$ give

$$2\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, \tilde{\mathcal{S}}_{2;0,0}|_{\sqrt{q}} = -4\left(\frac{\pi}{2}\right)^4 W_3\,, \tag{III.40}$$

including the $1/2^4$ present in (II.26). The $\tilde{\mathcal{S}}_{4;0,0}$ terms have two different kind of contributions: double trace and simple trace (see respectively first and second lines in (II.26)). In the spin structure $(0,0)$ the double trace always vanishes in $\tilde{\mathcal{S}}_{4;0,0}|_{\sqrt{q}}$ since

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, \frac{\sin(\pi\nu_{ij})}{\sin(\pi\nu_{kl})^2 \sin(\pi\nu_{ij})} = \int_{-\frac{1}{2}}^{\frac{1}{2}} \prod_{i=1}^{4} d\nu_i^1\, \frac{1}{\sin^2(\pi\nu_{kl})} = 0\,. \tag{III.41}$$

However the simple trace terms are treated in the same spirit as for the heterotic string. Only closed chains of sines contribute and are non zero only for specific ordering of the vertex operators. For instance,

$$-4\pi^4 \frac{\sin(\pi\nu_{ij})}{\sin(\pi\nu_{jk})\sin(\pi\nu_{kl})\sin(\pi\nu_{li})} \underset{\tau_2\to\infty}{\sim} -(2\pi)^4\,, \tag{III.42}$$

for the ordering $0 \le \omega_j < \omega_l < \omega_k < \omega_i \le 1$. Summing all of the contributions from $\tilde{\mathcal{S}}_{4;0,0}$ gives a total factor of $-\pi^4/16$, including the normalization in (II.26). We can now collect all the terms to get

$$A^{spin\ 2}_{(4,0)II} = \frac{1}{4}\left(\frac{\pi}{2}\right)^8 t_8 t_8 R^4\,(1 - 4W_3 + 2\,W^B)\,, \tag{III.43}$$

showing the equality with the heterotic expression

$$A^{spin\ 2}_{(4,0)het} = A^{spin\ 2}_{(4,0)II} \,. \tag{III.44}$$

Thanks to this equality for the spin 2 contribution and the one in (III.25) for the spin 1 contribution in the two $(4,0)$ models, from now we will use the notation $A^{spin\ s}_{(4,0)}$ with $s = 1, 2$.

The perturbative equality between these two $(4,0)$ models is not surprising. For a given number of vector multiplets n_v the heterotic and asymmetric type II construction lead to two string theory $(4,0)$ models related by S-duality, $S \to -1/S$, where S is the axion-dilaton complex scalar in the $\mathcal{N} = 4$ supergravity multiplet. The perturbative expansion in these two models is defined around different point in the $SU(1,1)/U(1)$ moduli space. The action of $\mathcal{N} = 4$ supersymmetry implies that the one-loop amplitudes between gravitons, which are neutral under the $U(1)$ R-symmetry, are the same in the strong and weak coupling regime.

2. *Comparing the* $(4,0)$ *and* $(2,2)$ *models*

In the case of the $(2,2)$ models, the contribution from the vector multiplets is given in (III.18). This is different from the expression in (III.8) for the $(4,0)$ as it can be easily seen using the supersymmetric Riemann identity in (III.5).

We now compare the spin $\frac{3}{2}$ contributions in the $(4,0)$ model in (III.16) and the $(2,2)$ model in (III.21). Performing the ν_i^1 integrations we can show that the effect from the spin $\frac{3}{2}$ multiplet running in the loop is the not the same in the two models but the difference is explained by the vector supermultiplet contribution

$$A^{spin\ \frac{3}{2}}_{(2,2)} = A^{spin\ \frac{3}{2}}_{(4,0)} - 2\,(A^{spin\ 1}_{(2,2)} - A^{spin\ 1}_{(4,0)}) \,. \tag{III.45}$$

We refer to section IV C 1 for the complete evaluation of the field theory one-loop amplitude where we show that the two spin 1 amplitude are different.

Therefore using the decomposition of the $\mathcal{N} = 4$ super-graviton contribution in the $(4,0)$ model in (III.14) and in the $(2,2)$ model in (III.20) we find that

$$A^{spin\ 2}_{(2,2)} = A^{spin\ 2}_{(4,0)} + 2\,(A^{spin\ 1}_{(2,2)} - A^{spin\ 1}_{(4,0)}) \,. \tag{III.46}$$

The difference between the spin 2 amplitudes in the two models is completely accounted for by the different vector multiplets contributions. This is not surprising since the string theory models are related by a U-duality exchanging the axion-dilaton scalar S of the gravity multiplet with a geometric modulus [21, 22, 31]. This transformation affects the coupling of the multiplet running in the loop explaining the difference between the two models. The $(2,2)$ construction leads to a different supergravity Lagrangian due to a different organization of the couplings to the vector fields, reflected in the different identification of the parameters of the moduli space in (II.1). We therefore see that constructing a supergravity theory as a double copy of two supersymmetric Yang-Mills theories, it is not enough to have the same number of real supercharges one needs to make sure that the theories are equivalent in perturbation.

IV. FIELD THEORY ONE-LOOP AMPLITUDES IN $\mathcal{N}=4$ SUPERGRAVITY

In this section we shall extract and compute the field theory limit $\alpha' \to 0$ of the one-loop string theory amplitudes studied previously. We show some relations between loop momentum power counting and the spin or supersymmetry of the multiplet running in the loop.

As was said, the region of the fundamental domain integration corresponding to the field theory amplitude is $\tau_2 \to \infty$, such that $t = \alpha' \tau_2$ is fixed. We then obtain a world-line integral of total proper time t.

The generic form of the field theory four-graviton one-loop amplitude for $\mathcal{N}=4$ supergravity with a spin s ($s = 1, \frac{3}{2}, 2$) $\mathcal{N}=4$ supermultiplet running is the loop is given by

$$M_X^{spin\ s} = \left(\frac{4}{\pi}\right)^4 \frac{\mu^{2\epsilon}}{\pi^{\frac{D}{2}}} \int_0^\infty \frac{dt}{t^{\frac{D-6}{2}}} \int_{\Delta_\omega} \prod_{i=1}^{3} d\omega_i \; e^{-\pi\, t\, Q(\omega)} \times A_X^{spin\ s}\,, \tag{IV.1}$$

where $D = 4-2\epsilon$ and X stands for the model, $X = (4,0)het$, $X = (4,0)II$ or $X = (2,2)$ and the respective amplitudes $A_X^{spin\ s}$ are given in sections III B and III C. We have set the overall normalization to unity.

The domain of integration $\Delta_\omega = [0,1]^3$ is decomposed into three regions $\Delta_w = \Delta_{(s,t)} \cup \Delta_{(s,u)} \cup \Delta_{(t,u)}$ given by the union of the (s,t), (s,u) and (t,u) domains. In the $\Delta_{(s,t)}$ domain the integration is performed over $0 \le \omega_1 \le \omega_2 \le \omega_3 \le 1$ where $Q(\omega) = -s\omega_1(\omega_3-\omega_2) - t(\omega_2-\omega_1)(1-\omega_3)$ with equivalent formulas obtained by permuting the external legs labels in the (t,u) and (s,u) regions (see [41] for details). We used that $s = -(k_1+k_2)^2$, $t = -(k_1+k_4)^2$ and $u = -(k_1+k_3)^2$ with our convention for the metric $(-+\cdots+)$.

We turn to the evaluation of the amplitudes. The main properties of the bosonic and fermionic propagators are provided in the appendix A 2. For this we will work with the helicity choice made in the previous section. This choice of polarization makes the intermediate steps easier as the expressions are explicitly gauge invariant.

A. Supersymmetry in the loop

Before evaluating the amplitude we discuss the action of supersymmetry on the structure of the one-loop amplitudes. An n-graviton amplitude in dimensional regularization with $D = 4 - 2\epsilon$ can generically be written in the following way:

$$M_{n;1} = \mu^{2\epsilon} \int \frac{d^D \ell}{(2\pi)^D} \frac{\mathfrak{N}(\epsilon_i, k_i; \ell)}{\ell^2 (\ell - k_1)^2 \cdots (\ell - \sum_{i=1}^{n-1} k_i)^2}, \tag{IV.2}$$

where the numerator is a polynomial in the loop momentum ℓ with coefficients depending on the external momenta k_i and polarization of the gravitons ϵ_i. For ℓ large this numerator behaves as $\mathfrak{N}(\epsilon_i, k_i; \ell) \sim \ell^{2n}$ in non-supersymmetric theories. In a $\mathcal{N}$ extended supergravity theory, supersymmetric cancellations improve this behaviour, which becomes $\ell^{2n-\mathcal{N}}$ where $\mathcal{N}$ is the number of four-dimensional supercharges:

$$\mathfrak{N}^{\mathcal{N}}(\epsilon_i, k_i; \ell) \sim \ell^{2n-\mathcal{N}} \qquad \text{for} \qquad |\ell| \to \infty \,. \tag{IV.3}$$

The dictionary between the Feynman integral presentation given in (IV.2) and the structure of the field theory limit of the string theory amplitude given in [42] tells that a first derivative of the bosonic propagator counts as one power of loop momentum $\partial\mathcal{P} \sim \ell$, $\partial^2\mathcal{P} \sim \ell^2$ and that fermionic propagators $S_{a,b} \sim 1$.

With this dictionary we find that in the $(4,0)$ model the integrand of the amplitudes have the following behaviour

$$\begin{aligned} A^{spin\ 1}_{(4,0)} &\sim \ell^4 \,, \\ A^{spin\ \frac{3}{2}}_{(4,0)} &\sim \ell^2 + \ell^4 \,, \\ A^{spin\ 2}_{(4,0)} &\sim 1 + \ell^2 + \ell^4 \,. \end{aligned} \tag{IV.4}$$

The spin 1 contribution to the four-graviton amplitude has four powers of loop momentum as required for an $\mathcal{N} = 4$ amplitude according (IV.3). The $\mathcal{N} = 4$ spin $\frac{3}{2}$ supermultiplet contribution can be decomposed into an $\mathcal{N} = 6$ spin $\frac{3}{2}$ supermultiplet term with two powers of loop momentum, and an $\mathcal{N}=4$ spin 1 supermultiplet contribution with four powers of loop momentum. The spin 2 contribution has an $\mathcal{N}=8$ spin 2 piece with no powers of loop momentum, an $\mathcal{N}=6$ spin $\frac{3}{2}$ piece with two powers of loop momentum and an $\mathcal{N}=4$ spin 1 piece with four powers of loop momentum.

For the $(2,2)$ construction we have the following behaviour

$$\begin{aligned} A^{spin\ 1}_{(2,2)} &\sim (\ell^2)^2\,, \\ A^{spin\ \frac{3}{2}}_{(2,2)} &\sim \ell^2 + (\ell^2)^2\,, \\ A^{spin\ 2}_{(2,2)} &\sim 1 + \ell^2 + (\ell^2)^2\,. \end{aligned} \tag{IV.5}$$

Although the superficial counting of the number of loop momenta is the same for each spin $s=1,\frac{3}{2},2$ in the two models, the precise dependence on the loop momentum differs in the two models, as indicated by the symbolic notation ℓ^4 and $(\ell^2)^2$. This is a manifestation of the model dependence on the vector multiplets contributions. As we will see below the order four terms in the loop momentum in the spin $\frac{3}{2}$ and spin 2 parts are due to the spin 1 part which will turn to be the only model dependent part.

At the level of the string amplitude, the multiplets running in the loop (spin 2 and spin 1) are naturally decomposed under the $\mathcal{N}=4$ supersymmetry group. However, at the level of the amplitudes in field theory it is convenient to group the various blocks according to the number of powers of loop momentum in the numerator

$$A^{spin\ s}_{\mathcal{N}=4s} \sim \ell^{4(2-s)}, \qquad s=1,\frac{3}{2},2\,, \tag{IV.6}$$

which is the same as organizing the terms according to the supersymmetry of the corresponding $\mathcal{N}=4s$ spin $s=1,\frac{3}{2},2$ supermultiplet. In this decomposition it is understood that for the two $\mathcal{N}=4$ models the dependence in the loop momenta is not identical.

From these blocks, one can reconstruct the contribution of the spin 2 $\mathcal{N}=4$ multiplet that we are concerned with using the following relations

$$M^{spin\ 2}_{X} = M^{spin\ 2}_{\mathcal{N}=8} - 4M^{spin\ \frac{3}{2}}_{\mathcal{N}=6} + 2M^{spin\ 1}_{X}\,, \tag{IV.7}$$

where the index X refers to the type of model, $(4,0)$ or $(2,2)$.

This supersymmetric decomposition of the one-loop amplitude reproduces the one given in [24–30] and the power counting obtained from a superspace construction of the amplitudes in [43]. Moreover, combined with the cancellations from gauge (diffeomorphism) invariance, it leads to the IR and UV behaviour discussed in [42, 44].

We shall come now to the evaluation of those integrals. We will see how the dependence on the loop momentum affects the structure of the one-loop amplitudes.

B. Model dependent part : $\mathcal{N}=4$ vector multiplet contribution

In this section we first compute the field theory amplitude with an $\mathcal{N}=4$ vector multiplet running in the loop for the two models. Then we provide an analysis of the IR and UV behaviour of these amplitudes.

1. Evaluation of the field theory amplitude

The contribution from the $\mathcal{N}=4$ spin 1 vector supermultiplets in the $(4,0)$ models is

$$M^{spin\ 1}_{(4,0)} = \left(\frac{4}{\pi}\right)^4 \frac{\mu^{2\epsilon}}{\pi^{\frac{D}{2}}} \int_0^\infty \frac{dt}{t^{\frac{D-6}{2}}} \int_{\Delta_\omega} d^3\omega\ e^{-\pi\, t\, Q(\omega)} \times A^{spin\ 1}_{(4,0)}\,, \qquad \text{(IV.8)}$$

where $A^{spin\ 1}_{(4,0)}$ is given in (III.22) for instance and Q defined in (A.17). Integrating over the proper time t and setting $D=4-2\epsilon$, the amplitude reads

$$M^{spin\ 1}_{(4,0)} = t_8t_8R^4 \int_{\Delta_\omega} d^3\omega \big[\Gamma\left(1+\epsilon\right)\, Q^{-1-\epsilon}\, W_2 + \Gamma\left(2+\epsilon\right)\, Q^{-2-\epsilon}\, W_1\big]\ . \qquad \text{(IV.9)}$$

The quantities W_1 and W_2 are given in (III.38), they have the following form in terms of the variables ω_i:

$$W_1 = \frac{1}{8}\,(\omega_2-\omega_3)(\mathrm{sign}(\omega_1-\omega_2)+2\omega_2-1)(\mathrm{sign}(\omega_2-\omega_1)+2\omega_1-1)(\mathrm{sign}(\omega_3-\omega_2)+2\omega_2-1)$$

$$W_2 = \frac{1}{16}\,\frac{1}{u}\,(2\omega_2-1+\mathrm{sign}(\omega_3-\omega_2))(2\omega_2-1+\mathrm{sign}(\omega_1-\omega_2))\,(1-\delta(\omega_{24}))\,. \qquad \text{(IV.10)}$$

Using the dictionary given in [42], we recognize in the first term in (IV.9) a six dimensional scalar box integral, and in the second term four dimensional scalar bubble integrals. Evaluating the integrals using standard techniques, we find:[5]

$$M^{spin\ 1}_{(4,0)} = \frac{t_8t_8R^4}{2s^4}\left(s^2 - s(u-t)\log\left(\frac{-t}{-u}\right) - tu(\log^2\left(\frac{-t}{-u}\right)+\pi^2)\right)\ . \qquad \text{(IV.11)}$$

The crossing symmetry of the amplitudes has been broken by our choice of helicity configurations. The same comment applies to all the field theory amplitudes evaluated in this paper. This result matches the one derived in [24–29] and in particular [30, eq. (3.20)].

---

[5] The analytic continuation in the complex energy plane corresponds to the $+i\varepsilon$ prescription for the Feynman propagators $1/(\ell^2-m^2+i\varepsilon)$. We are using the notation that $\log(-s)=\log(-s-i\varepsilon)$ and that $\log(-s/-t):=\log((-s-i\varepsilon)/(-t-i\varepsilon))$.

Now we turn to the amplitude in the $(2,2)$ models:

$$M_{(2,2)}^{spin\ 1} = \left(\frac{4}{\pi}\right)^4 \frac{\mu^{2\epsilon}}{\pi^{\frac{D}{2}}} \int_0^\infty \frac{dt}{t^{\frac{D-6}{2}}} \int_{\Delta_\omega} d^3\omega\ e^{-\pi\, t\, Q(\omega)} \times A_{(2,2)}^{spin\ 1}\,, \tag{IV.12}$$

where $A_{(2,2)}^{spin\ 1}$ is defined in (III.18). After integrating over the proper time t, one gets

$$M_{(2,2)}^{spin\ 1} = \frac{t_8 t_8 R^4}{4} \int_{\Delta_\omega} d^3\omega\, \Gamma\left(2+\epsilon\right)\, Q^{-2-\epsilon} \left(W_3\right)^2 , \tag{IV.13}$$

where W_3 defined in (III.34), is given in terms of the ω_i variables by

$$\begin{aligned} W_3 = -\frac{1}{8}\,&(\mathrm{sign}(\omega_1-\omega_2)+2\omega_2-1)(\mathrm{sign}(\omega_2-\omega_1)+2\omega_1-1) \\ &\qquad\qquad\qquad -\frac{1}{8}\,(\mathrm{sign}(\omega_3-\omega_2)+2\omega_2-1)(\omega_3-\omega_2)\,. \end{aligned} \tag{IV.14}$$

Performing all the integrations leads to

$$\begin{aligned} M_{(2,2)}^{spin\ 1} = \frac{t_8 t_8 R^4}{4}\Bigg[& \\ \frac{1}{u^2}\left(\log^2\left(\frac{-s}{-t}\right)+\pi^2\right) &+ \frac{2u(u-5t)}{s^4}\left(\log^2\left(\frac{-t}{-u}\right)+\pi^2\right) - \frac{2u}{t^3}\left(\log^2\left(\frac{-u}{-s}\right)+\pi^2\right) \\ &+\frac{2}{su}\log\left(\frac{-s}{-t}\right) - \frac{8(2u-t)}{s^3}\log\left(\frac{-t}{-u}\right) - \frac{2(s-u)}{st^2}\log\left(\frac{-u}{-s}\right) \\ &+\frac{2(s+6t)}{ts^2}\Bigg]\,. \end{aligned} \tag{IV.15}$$

This expression is the linear combination of six dimensional scalar boxes, four dimensional scalar bubbles and a rational piece in agreement with the dictionary given in [42].

It is now clear that the vector multiplet contribution to the amplitude differs in the two theories, $(4,0)$ and $(2,2)$. It would be interesting to see if this expression can be derived using the unitarity methods of [24–29] and the double-copy construction of [30].

2. *IR and UV behaviour*

The graviton amplitudes with vector multiplets running in the loop in (IV.11) and (IV.15) are free of UV and IR divergences. The absence of IR divergence is expected, since no spin 2 state is running in the loop. The IR divergence occurs only when a graviton is exchanged between two soft graviton legs (see figure 1). This fact has already been noticed in [25].

This behaviour is easily understood by considering the soft graviton limit of the coupling between the graviton and a spin $s \neq 2$ state. It occurs through the stress-energy tensor



FIG. 1. Contribution to the IR divergences when two external graviton (double wavy-lines) become soft. If a graviton is exchanged as in (a) the amplitude presents an IR divergence. No IR divergences are found when another massless state of spin different from two is exchanged as in figure (b).

$V^{\mu\nu}(k,p) = T^{\mu\nu}(p-k,p)$ where k and p are respectively the momentum of the graviton and of the exchanged state. In the soft graviton limit the vertex behaves as $V^{\mu\nu}(p-k,p) \sim -k^\mu p^\nu$ for $p^\mu \sim 0$, and the amplitude behaves in the soft limit as

$$\int_{\ell\sim 0} \frac{d^4\ell}{\ell^2(\ell\cdot k_1)(\ell\cdot k_2)}\, T_{\mu\nu}(\ell-k_1,\ell)T^{\mu\nu}(\ell,\ell+k_2) \sim (k_1\cdot k_2)\int_{\ell\sim 0} \frac{d^4\ell}{\ell^2(\ell\cdot k_1)(\ell\cdot k_2)}\,\ell^2\,, \quad \text{(IV.16)}$$

which is finite for small values of the loop momentum $\ell \sim 0$. In the soft graviton limit, the three graviton vertex behaves as $V^{\mu\nu}(k,p) \sim k^\mu k^\nu$ and the amplitude has a logarithmic divergence at $\ell \sim 0$

$$(k_1\cdot k_2)^2 \int_{\ell\sim 0} \frac{d^4\ell}{\ell^2(\ell\cdot k_1)(\ell\cdot k_2)} = \infty\,. \quad \text{(IV.17)}$$

The absence of UV divergence is due to the fact that the R^2 one-loop counter-term is the Gauss-Bonnet term. It vanishes in the four-point amplitude since it is a total derivative [45].

C. Model independent part

In this section we compute the field theory amplitudes with an $\mathcal{N} = 8$ supergraviton and an $\mathcal{N} = 6$ spin $\frac{3}{2}$ supermultiplet running in the loop. These quantities are model independent.

1. The $\mathcal{N} = 6$ spin $\frac{3}{2}$ supermultiplet contribution

The $\mathcal{N} = 4$ spin $\frac{3}{2}$ supermultiplet contribution is different in the two $(4,0)$ and $(2,2)$ constructions of the $\mathcal{N} = 4$ supergravity models. As shown in equation (III.45), this is accounted for by the contribution of the vector multiplets. However, we exhibit an $\mathcal{N} = 6$

spin $\frac{3}{2}$ supermultiplet model independent piece by adding two $\mathcal{N} = 4$ vector multiplets contributions to the one of an $\mathcal{N} = 4$ spin $\frac{3}{2}$ supermultiplet

$$M^{spin\ \frac{3}{2}}_{\mathcal{N}=6} = M^{spin\ \frac{3}{2}}_{X} + 2M^{spin\ 1}_{X} \,. \tag{IV.18}$$

The amplitude with an $\mathcal{N} = 6$ spin $\frac{3}{2}$ multiplet running in the loop is

$$M^{spin\ \frac{3}{2}}_{\mathcal{N}=6} = -\frac{t_8 t_8 R^4}{8} \int_{\Delta_\omega} d^3\omega\, \Gamma\left(2+\epsilon\right)\, W_3\, Q^{-2-\epsilon}, \tag{IV.19}$$

where W_3 is given in (IV.14). The integral is equal to the six dimensional scalar box integral given in [30, eq. (3.16)] up to $O(\epsilon)$ terms. We evaluate it, and get

$$M^{spin\ \frac{3}{2}}_{\mathcal{N}=6} = -\frac{t_8 t_8 R^4}{2s^2} \left(\log^2\left(\frac{-t}{-u}\right) + \pi^2\right) \,. \tag{IV.20}$$

This result is UV finite as expected from the superficial power counting of loop momentum in the numerator of the amplitude given in (IV.4). It is free of IR divergences because no graviton state is running in the loop (see the previous section). It matches the one derived in [24–29] and in particular [30, eq. (3.17)].

2. The $\mathcal{N} = 8$ spin 2 supermultiplet contribution

We now turn to the $\mathcal{N} = 8$ spin 2 supermultiplet contribution in (IV.7). It has already been evaluated in [43, 46] and writes:

$$M^{spin\ 2}_{\mathcal{N}=8} = \frac{t_8 t_8 R^4}{4} \int_{\Delta_\omega} d^3\omega\, \Gamma\left(2+\epsilon\right)\, Q^{-2-\epsilon} \,. \tag{IV.21}$$

Performing the integrations we have

$$\begin{aligned} M^{spin\ 2}_{\mathcal{N}=8} = \frac{t_8 t_8 R^4}{4} \Bigg[\frac{2}{\epsilon} \left(\frac{\log\left(\frac{-t}{\mu^2}\right)}{su} + \frac{\log\left(\frac{-s}{\mu^2}\right)}{tu} + \frac{\log\left(\frac{-u}{\mu^2}\right)}{st} \right) + \\ + 2 \left(\frac{\log\left(\frac{-s}{\mu^2}\right)\log\left(\frac{-t}{\mu^2}\right)}{st} + \frac{\log\left(\frac{-t}{\mu^2}\right)\log\left(\frac{-u}{\mu^2}\right)}{tu} + \frac{\log\left(\frac{-u}{\mu^2}\right)\log\left(\frac{-s}{\mu^2}\right)}{us} \right) \Bigg] \,, \end{aligned} \tag{IV.22}$$

where μ^2 is an IR mass scale. This amplitudes carries an ϵ pole signaling the IR divergence due to the graviton running in the loop.

Now we have all the blocks entering the expression for the $\mathcal{N} = 4$ pure gravity amplitude in (IV.7).

V. CONCLUSION

In this work we have evaluated the four-graviton amplitude at one-loop in $\mathcal{N}=4$ supergravity in four dimensions from the field theory limit of string theory construction. The string theory approach provides $(4,0)$ models where all supersymmetry arise from the left-moving sector of the theory, and $(2,2)$ models where supersymmetry arises from both the left and right moving sectors of the theory.

For each model this gives a pure $\mathcal{N}=4$ supergravity four-graviton one-loop amplitude, defined by the subtraction of the vector multiplet contributions. We have seen that, except when gravitons are running in the loop, the one-loop amplitudes are free of IR divergences. In addition, all the amplitudes are UV finite because the R^2 candidate counter-term vanishes for these amplitudes. Amplitudes with external vector multiplets states are expected to be UV divergent [47].

In the $(4,0)$ setup, our results match the ones obtained with the unitarity based method in [24–29] and the double-copy approach of [30]. This shows that, for a supersymmetric theory, when an explicit string theory representation is available one can easily obtain the corresponding field theory amplitudes.

The string theory construction shows that in the $(2,2)$ models the one-loop amplitude take a different form. This difference can be traced to the different nature of the couplings of the vector-multiplet in the two theory as indicated by the relation between the two amplitudes in (III.46). In this representation, the one-loop amplitudes can be organized as a linear combination of $\mathcal{N}=4s$ spin $s=1,\frac{3}{2},2$ contributions, with a simple power counting dependence on the loop momentum $\ell^{4(2-s)}$.

As we have said, the one-loop vector multiplet amplitudes in (IV.11) and (IV.15) extracted from the two string theory constructions differ. This is expected as these amplitudes are extracted from string theory constructions evaluated around two $\mathcal{N}=4$ vacua with a different identification of the scalar fields moduli in the gravity and matter multiplet. It would be interesting to reproduce these results using a double-copy construction from $\mathcal{N}=2$ super-Yang-Mills amplitudes.

ACKNOWLEDGEMENTS

We would like to thank Ashoke Sen for discussions and in particular Arniban Basu for having pointed to us the construction by Dabholkar and Harvey in [20]. We would like to thank Guillaume Bossard, Renata Kallosh and Warren Siegel for comments on a previous version of this paper.

Appendix A: World-sheet CFT : chiral blocks, propagators.

In this appendix we collect various results about the conformal blocks, fermionic and bosonic propagators at genus one, and their q-expansions.

1. Bosonic and fermionic chiral blocks

▷ The genus one theta functions are defined to be

$$\theta \begin{bmatrix} a \\ b \end{bmatrix} (z|\tau) = \sum_{n\in\mathbb{Z}} q^{\frac{1}{2}(n+\frac{a}{2})^2} \, e^{2i\pi(z+\frac{b}{2})(n+\frac{a}{2})} \,, \tag{A.1}$$

and Dedekind eta function:

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1-q^n) \,, \tag{A.2}$$

where $q = \exp(2i\pi\tau)$. Those functions have the following $q \to 0$ behaviour:

$$\begin{aligned} &\theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (0,\tau) = 0 \,; \quad \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0,\tau) = -2q^{1/8} + O(q) \,; \quad \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0,\tau) = \frac{1}{\sqrt{q}} + 2 + O(q) \,; \\ &\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix} (0,\tau) = \frac{1}{\sqrt{q}} - 2 + O(q) \,; \quad \eta(\tau) = q^{1/24} + O(q) \,. \end{aligned} \tag{A.3}$$

▷ The partition function of eight world-sheet fermion in the (a,b)-spin structure, $\Psi(z+1) = -(-1)^{2a}\Psi(z)$ and $\Psi(z+\tau) = -(-1)^{2b}\Psi(z)$, and eight chiral bosons is

$$Z_{a,b}(\tau) \equiv \frac{\theta \left[\begin{smallmatrix} a \\ b \end{smallmatrix}\right] (0|\tau)^4}{\eta^{12}(\tau)} \,, \tag{A.4}$$

it has the following behaviour for $q \to 0$

$$\begin{aligned} Z_{1,1} &= 0 \,, \\ Z_{1,0} &= 16 + 16^2 q + o(q^2) \,, \\ Z_{0,0} &= \frac{1}{\sqrt{q}} + 8 + o(\sqrt{q}) \,, \\ Z_{0,1} &= \frac{1}{\sqrt{q}} - 8 + o(\sqrt{q}) \,. \end{aligned} \tag{A.5}$$

▷ The partition function of the twisted (X,Ψ) system in the (a,b)-spin structure is

$$\begin{aligned} X(z+1) &= (-1)^{2h} X(z); \qquad \Psi(z+1) = -(-1)^{2a+2h}\Psi(z) \,, \\ X(z+\tau) &= (-1)^{2g} X(z); \qquad \Psi(z+\tau) = -(-1)^{2b+2g}\Psi(z) \,. \end{aligned} \tag{A.6}$$

The twisted chiral blocks for a real boson are

$$\mathcal{Z}^{h,g}[X] = \left(i e^{-i\pi g} q^{-h^2/2} \frac{\eta(\tau)}{\theta\begin{bmatrix}1+h\\1+g\end{bmatrix}} \right)^{1/2} . \tag{A.7}$$

The twisted chiral blocks for a Majorana or Weyl fermion are

$$\mathcal{Z}^{h,g}_{a,b}[\Psi] = \left(e^{-i\pi\, a(g+b)/2} q^{h^2/2} \frac{\theta\begin{bmatrix}a+h\\b+g\end{bmatrix}}{\eta(\tau)} \right)^{1/2} . \tag{A.8}$$

The total partition function is given by

$$\mathcal{Z}^{h,g}_{a,b}[(x,\Psi)] = \mathcal{Z}^{h,g}[X]\mathcal{Z}^{h,g}_{a,b}[\Psi] = e^{i\frac{\pi}{4}(1+2g+a(g+b))} \sqrt{\frac{\theta\begin{bmatrix}a+h\\b+g\end{bmatrix}}{\theta\begin{bmatrix}1+h\\1+g\end{bmatrix}}} . \tag{A.9}$$

2. Bosonic and fermionic propagators

a. Bosonic propagators

Our convention for the bosonic propagator is

$$\langle x^\mu(\nu) x^\nu(0)\rangle_{one-loop} = 2\alpha'\, \eta^{\mu\nu}\, \mathcal{P}(\nu|\tau) , \tag{A.10}$$

with

$$\begin{aligned} \mathcal{P}(\nu|\tau) &= -\frac{1}{4}\ln\left|\frac{\theta\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right](\nu|\tau)}{\partial_\nu\theta\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right](0|\tau)}\right|^2 + \frac{\pi\nu_2^2}{2\tau_2} + C(\tau) \\ &= \frac{\pi\nu_2^2}{2\tau_2} - \frac{1}{4}\ln\left|\frac{\sin(\pi\nu)}{\pi}\right|^2 - \sum_{m\geq 1}\left(\frac{q^m}{1-q^m}\frac{\sin^2(m\pi\nu)}{m} + c.c.\right) + C(\tau), \end{aligned} \tag{A.11}$$

where $C(\tau)$ is a contribution of the zero modes (see e.g. [41]) that anyway drops out of the string amplitude because of momentum conservation so we will forget it in the following.

We have as well the expansions

$$\begin{aligned} \partial_\nu\mathcal{P}(\nu|\tau) &= \frac{\pi}{2i}\,\frac{\nu_2}{\tau_2} - \frac{\pi}{4}\frac{1}{\tan(\pi\nu)} - \pi\, q\, \sin(2\pi\nu) + O(q) , \\ \partial_\nu^2\mathcal{P}(\nu|\tau) &= -\frac{\pi}{4\tau_2} + \frac{\pi^2}{4}\frac{1}{\sin^2(\pi\nu)} - 2\pi^2\, q\cos(2\pi\nu) + O(q) . \end{aligned} \tag{A.12}$$

leading to the following Fourier expansion with respect to ν_1

$$\partial_\nu \mathcal{P}(\nu|\tau) = \frac{\pi}{4i}\,(2\nu_2 - \mathrm{sign}(\nu_2)) + i\frac{\pi}{4}\,\mathrm{sign}(\nu_2)\sum_{m\neq 0} e^{2i\pi m \mathrm{sign}(\nu_2)\nu} - \pi\, q\, \sin(2\pi\nu) + O(q)\,,$$
$$\partial_\nu^2 \mathcal{P}(\nu|\tau) = \frac{\pi}{4\tau_2}\,(\tau_2\delta(\nu_2) - 1) - \pi^2 \sum_{m\geq 1} m e^{2i\pi m \mathrm{sign}(\nu_2)\nu} - 2\pi^2\, q\, \cos(2\pi\nu) + O(q)\,, \quad \text{(A.13)}$$

Setting $\nu = \nu_1 + i\tau_2\omega$ we can rewrite these expressions in a form relevant for the field theory limit $\tau_2 \to \infty$ with $t = \alpha'\tau_2$ =fixed. The bosonic propagator can be decomposed in an asymptotic value for $\tau_2 \to \infty$ (the field theory limit) and corrections originating from massive string modes

$$\mathcal{P}(\nu|\tau) = -\frac{\pi\, t}{2\alpha'}\, P(\omega) + \delta P(\nu) - q\, \sin^2(\pi\nu) - \bar{q}\, \sin^2(\pi\bar{\nu}) + O(q^2)\,, \quad \text{(A.14)}$$

and

$$P(\omega) = \omega^2 - |\omega|; \qquad \delta P(\nu) = \sum_{m\neq 0} \frac{1}{4|m|}\, e^{2i\pi m\nu_1 - 2\pi|m\nu_2|}\,. \quad \text{(A.15)}$$

The contribution δP corresponds to the effect of massive string states propagating between two external massless states. The quantity $\mathcal{Q}$ defined in (II.4) writes in this limit

$$\mathcal{Q} = -t\pi Q(\omega) + \alpha'\delta Q - 2\pi\alpha' \sum_{1\leq i<j\leq 4} k_i\cdot k_j\, (q \sin^2(\pi\nu_{ij}) + \bar{q}\sin^2(\pi\bar{\nu}_{ij})) + O(q^2)\,, \quad \text{(A.16)}$$

where

$$Q(\omega) = \sum_{1\leq i<j\leq 4} k_i\cdot k_j\, P(\omega_{ij})\,, \qquad \delta Q = 2\sum_{1\leq i<j\leq 4} k_i\cdot k_j \delta P(\nu_{ij})\,. \quad \text{(A.17)}$$

b. *Fermionic propagators*

Our normalization for the fermionic propagators in the (a,b) spin structure is given by

$$\langle \psi^\mu(z)\psi^\nu(0)\rangle_{one-loop} = \frac{\alpha'}{2}\, S_{a,b}(z|\tau)\,. \quad \text{(A.18)}$$

▷ In the even spin structure fermionic propagators are

$$S_{a,b}(z|\tau) = \frac{\theta\left[\begin{smallmatrix}a\\b\end{smallmatrix}\right](z|\tau)}{\theta\left[\begin{smallmatrix}a\\b\end{smallmatrix}\right](0|\tau)}\, \frac{\partial_z\theta\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right](0|\tau)}{\theta\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right](z|\tau)}\,. \quad \text{(A.19)}$$

The odd spin structure propagator is

$$S_{1,1}(z|\tau) = \frac{\partial_z\theta\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right](z|\tau)}{\theta\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right](z|\tau)}\,, \quad \text{(A.20)}$$

the fermionic propagator orthogonal to the zero modes is

$$\tilde{S}_{1,1}(z|\tau) = S_{1,1}(z|\tau) - 2i\pi\frac{z_2}{\tau_2} = -4\partial_z\mathcal{P}(z|\tau)\,. \tag{A.21}$$

The fermionic propagators have the following q-expansion representation [48]

$$\begin{aligned}
S_{1,1}(z|\tau) &= \frac{\pi}{\tan(\pi z)} + 4\pi\sum_{n=1}^{\infty}\frac{q^n}{1-q^n}\,\sin(2n\pi z)\,,\\
S_{1,0}(z|\tau) &= \frac{\pi}{\tan(\pi z)} - 4\pi\sum_{n=1}^{\infty}\frac{q^n}{1+q^n}\,\sin(2n\pi z)\,,\\
S_{0,0}(z|\tau) &= \frac{\pi}{\sin(\pi z)} - 4\pi\sum_{n=1}^{\infty}\frac{q^{n-\frac{1}{2}}}{1+q^{n-\frac{1}{2}}}\,\sin((2n-1)\pi z)\,,\\
S_{0,1}(z|\tau) &= \frac{\pi}{\sin(\pi z)} + 4\pi\sum_{n=1}^{\infty}\frac{q^{n-\frac{1}{2}}}{1-q^{n-\frac{1}{2}}}\,\sin((2n-1)\pi z)\,.
\end{aligned} \tag{A.22}$$

▷ Riemann supersymmetric identies written in the text (II.27) derive from the following Riemann relation relation:

$$\sum_{a,b=0,1}(-1)^{a+b+ab}\prod_{i=1}^{4}\theta\begin{bmatrix}a\\b\end{bmatrix}(v_i) = -2\prod_{i=1}^{4}\theta_1(v_i')\,, \tag{A.23}$$

with $v_1' = \frac{1}{2}(-v_1+v_2+v_3+v_4)$ $v_2' = \frac{1}{2}(v_1-v_2+v_3+v_4)$ $v_3' = \frac{1}{2}(v_1+v_2-v_3+v_4)$ $v_4' = \frac{1}{2}(v_1+v_2+v_3-v_4)$. This identity can be written, in the form used in the main text, as vanishing identities

$$\sum_{\substack{a,b=0,1\\ab=0}}(-1)^{a+b+ab}Z_{a,b}(\tau) = 0\,, \tag{A.24}$$

$$\sum_{\substack{a,b=0,1\\ab=0}}(-1)^{a+b+ab}Z_{a,b}(\tau)\prod_{r=1}^{n}S_{a,b}(z) = 0 \quad 1\leq n\leq 3\,, \tag{A.25}$$

and the first non-vanishing one

$$\sum_{\substack{a,b=0,1\\ab=0}}(-1)^{a+b+ab}Z_{a,b}(\tau)\prod_{i=1}^{4}S_{a,b}(z_i|\tau) = -(2\pi)^4\,. \tag{A.26}$$

with $z_1+\cdots+z_4=0$ and where we used that $\partial_z\theta\left[\begin{smallmatrix}1\\1\end{smallmatrix}\right](0|\tau) = 2\pi\eta^3(\tau)$.

c. *q-expansion*

The q-expansions of the fermionic propagators in the even spin structure are given by

$$S_{1,0}(z|\tau) = \frac{\pi}{\tan(\pi z)} - 4\pi q\sin(2\pi z) + O(q^2)\,,$$

$$
\begin{aligned}
S_{0,0}(z|\tau) &= \frac{\pi}{\sin(\pi z)} - 4\pi\sqrt{q}\,\sin(\pi z) + O(q)\,, \\
S_{0,1}(z|\tau) &= \frac{\pi}{\sin(\pi z)} + 4\pi\sqrt{q}\,\sin(\pi z) + O(q)\,.
\end{aligned}
\tag{A.27}
$$

Setting $\mathcal{S}^n_{a,b} = \prod_{i=1}^n S_{a,b}(z_i|\tau)$ we have the following expansion

$$
\begin{aligned}
S^n_{1,0} &= \prod_{i=1}^n \pi\cot(\pi z_i)\,\left(1 - 8q\sum_{i=1}^n \sin^2(\pi z_i)\right) + o(q^2)\,, \\
S^n_{0,0} &= \prod_{i=1}^n \pi(\sin(\pi z_i))^{-1}\,\left(1 - 4q\sum_{i=1}^n \sin^2(\pi z_i)\right) + o(q^2)\,, \\
S^n_{0,1} &= \prod_{i=1}^n \pi(\sin(\pi z_i))^{-1}\,\left(1 + 4q\sum_{i=1}^n \sin^2(\pi z_i)\right) + o(q^2)\,.
\end{aligned}
\tag{A.28}
$$

Applying these identities with $n = 2$ and $n = 4$ we derive the following relations between the correlators $\mathcal{W}^F_{a,b}$ defined in (II.25)

$$
\mathcal{W}^F_{0,0}|_{q^0} = \mathcal{W}^F_{0,1}|_{q^0}; \qquad \mathcal{W}^F_{0,0}|_{\sqrt{q}} = -\mathcal{W}^F_{0,1}|_{\sqrt{q}}\,. \tag{A.29}
$$

Using the q-expansion of the bosonic propagator, it is not difficult to realize that $\mathcal{W}^B|_{\sqrt{q}} = 0$, and we can promote the previous relation to the full correlator $\mathcal{W}_{a,b}$ defined in (II.23)

$$
\mathcal{W}_{0,0}|_{q^0} = \mathcal{W}_{0,1}|_{q^0}; \qquad \mathcal{W}_{0,0}|_{\sqrt{q}} = -\mathcal{W}_{0,1}|_{\sqrt{q}}\,. \tag{A.30}
$$

Other useful relations are between the q-expansion of the derivative bosonic propagator $\partial\mathcal{P}$ and the fermionic propagator $S_{1,0}$

$$
\begin{aligned}
\partial_\nu \mathcal{P}(\nu|\tau)|_{q^0} - \frac{\pi\nu_2}{2i\tau_2} &= -\frac{1}{4}S_{1,0}(\nu|\tau)|_{q^0} \\
\partial_\nu \mathcal{P}(\nu|\tau)|_{q^1} &= +\frac{1}{4}S_{1,0}(\nu|\tau)|_{q}\,.
\end{aligned}
\tag{A.31}
$$

3. Congruence subgroups of $SL(2,\mathbb{Z})$

We denote by $SL(2,\mathbb{Z})$ the group of 2×2 matrix with integers entries of determinant 1. For any N integers we have the following subgroups of $SL(2,\mathbb{Z})$

$$\begin{aligned}\Gamma_0(N) &= \left\{\begin{pmatrix} a & b\\ c & d\end{pmatrix}\in SL(2,\mathbb{Z})|\begin{pmatrix} a & b\\ c & d\end{pmatrix}=\begin{pmatrix} * & *\\ 0 & *\end{pmatrix}\quad \mathrm{mod}\ N\right\}\,,\\ \Gamma_1(N) &= \left\{\begin{pmatrix} a & b\\ c & d\end{pmatrix}\in SL(2,\mathbb{Z})|\begin{pmatrix} a & b\\ c & d\end{pmatrix}=\begin{pmatrix} 1 & *\\ 0 & 1\end{pmatrix}\quad \mathrm{mod}\ N\right\}\,,\\ \Gamma(N) &= \left\{\begin{pmatrix} a & b\\ c & d\end{pmatrix}\in SL(2,\mathbb{Z})|\begin{pmatrix} a & b\\ c & d\end{pmatrix}=\begin{pmatrix} 1 & 0\\ 0 & 1\end{pmatrix}\quad \mathrm{mod}\ N\right\}\,.\end{aligned}\tag{A.32}$$

They satisfy the properties that $\Gamma(N)\subset\Gamma_1(N)\subset\Gamma_0(N)\subset SL(2,\mathbb{Z})$.

Appendix B: Chiral blocks for the type II orbifolds

We recall some essential facts from the construction of [23]. The shifted T^2 lattice sum writes

$$\Gamma^w_{(2,2)}\begin{bmatrix} h\\ g\end{bmatrix} := \sum_{P_L,p_R\in\Gamma_{(2,2)}+w\frac{h}{2}} e^{i\pi g l\cdot w} q^{\frac{P_L^2}{2}}\bar{q}^{\frac{P_R^2}{2}}\,,\tag{B.1}$$

where $\ell\cdot w=m_I b^I+a_I n^I$ where the and the shift vector $w=(a_I,b^I)$ is such that $w^2=2a\cdot b=0$ and

$$\begin{aligned}P_L^2 &= \frac{|U(m_1+a_1\frac{h}{2})-(m_2+a_2\frac{h}{2})+T(n^1+b^1\frac{h}{2})+TU(n^2+b^2\frac{h}{2})|^2}{2T_2U_2}\,,\\ P_L^2-P_R^2 &= 2(m_I+a_I\frac{h}{2})(n^I+b^I\frac{h}{2})\,.\end{aligned}\tag{B.2}$$

T and U are the moduli of the T^2. We recall the full expressions for the orbifold blocks :

$$\mathcal{Z}^{(22);h,g}_{a,b} := \begin{cases}\mathcal{Z}_{a,b}=(\mathrm{II}.20) & (h,g)=(0,0)\\ 4(-1)^{(a+h)g}\left(\frac{\theta\left[{a\atop b}\right](0|\tau)\theta\left[{a+h\atop b+g}\right](0|\tau)}{\eta(\tau)^3\theta\left[{1+h\atop 1+g}\right]}\right)^2\times\Gamma_{(2,2)}(T,U) & (h,g)\neq(0,0)\,,\end{cases}\tag{B.3}$$

$$\mathcal{Z}^{(14);h,g}_{a,b} = \frac{1}{2}\sum_{h',g'=0}^{1}\mathcal{Z}^{h,g}_{a,b}\begin{bmatrix} h'\\ g'\end{bmatrix}\Gamma^w_{(2,2)}\begin{bmatrix} h'\\ g'\end{bmatrix}\,,\tag{B.4}$$

$$\mathcal{Z}^{(10);h,g}_{a,b} = \frac{1}{2}\sum_{h_1,g_1=0}^{1}\frac{1}{2}\sum_{h_2,g_2=0}^{1} \mathcal{Z}^{h,g}_{a,b}\begin{bmatrix}h;h_1,h_2\\ g;g_1,g_2\end{bmatrix}\Gamma^{w_1,w_2}_{(2,2)}\begin{bmatrix}h_1,h_2\\ g_1,g_2\end{bmatrix}\,, \forall h,g\,. \tag{B.5}$$

For th $n_v = 6$ model, the orbifold acts differently and we get

$$\mathcal{Z}^{(6);h,g}_{a,b} = \frac{1}{2}\sum_{h',g'=0}^{1}(-1)^{hg'+gh'}\,\mathcal{Z}^{h,g}_{a,b}\Gamma^{w}_{(2,2)}\begin{bmatrix}h'\\ g'\end{bmatrix}\,. \tag{B.6}$$

In the previous expressions, the crucial point is that the shifted lattice sums $\Gamma^{w}_{(2,2)}\begin{bmatrix}h'\\ g'\end{bmatrix}$ act as projectors on their untwisted $h' = 0$ sector, while the g' sector is left free. We recall now the diagonal properties of the orbifold action (see [23] again) on the lattice sums:

$$\Gamma^{w_1,w_2}_{(2,2)}\begin{bmatrix}h,0\\ g,0\end{bmatrix} = \Gamma^{w_1}_{(2,2)}\begin{bmatrix}h\\ g\end{bmatrix}\,,\ \Gamma^{w_1,w_2}_{(2,2)}\begin{bmatrix}0,h\\ 0,g\end{bmatrix} = \Gamma^{w_2}_{(2,2)}\begin{bmatrix}h\\ g\end{bmatrix}\,,\ \Gamma^{w_1,w_2}_{(2,2)}\begin{bmatrix}h,h\\ g,g\end{bmatrix} = \Gamma^{w_{12}}_{(2,2)}\begin{bmatrix}h\\ g\end{bmatrix}\,, \tag{B.7}$$

The four dimensional blocks $\mathcal{Z}^{h,g}_{a,b}$ have the following properties : $\mathcal{Z}^{h,g}_{a,b}\begin{bmatrix}0\\ 0\end{bmatrix} = \mathcal{Z}^{h,g}_{a,b}\begin{bmatrix}h\\ g\end{bmatrix} = \mathcal{Z}^{h,g}_{a,b}$ (ordinary twist); $\mathcal{Z}^{0,0}_{a,b}\begin{bmatrix}h\\ g\end{bmatrix}$ is a $(4,4)$ lattice sum with one shifted momentum thus is projects out the $h = 0$ sector. Equivalent properties stand as well for the $n_v = 10$ model.

One has then in the field theory limit

$$\begin{aligned}
\mathcal{Z}^{(14);h,g}_{a,b} &\in \{\mathcal{Z}^{0,0}_{a,b},\ \mathcal{Z}^{0,1}_{a,b},\ \frac{1}{2}\mathcal{Z}^{1,0}_{a,b},\ \frac{1}{2}\mathcal{Z}^{1,1}_{a,b}\}\,,\\
\mathcal{Z}^{(10);h,g}_{a,b} &\in \{\mathcal{Z}^{0,0}_{a,b},\ \mathcal{Z}^{0,1}_{a,b},\ \frac{1}{4}\mathcal{Z}^{1,0}_{a,b},\ \frac{1}{4}\mathcal{Z}^{1,1}_{a,b}\}\,,\\
\mathcal{Z}^{(6);h,g}_{a,b} &\in \{\mathcal{Z}^{0,0}_{a,b},\ \mathcal{Z}^{0,1}_{a,b},\ 0,\ 0\}\,,
\end{aligned} \tag{B.8}$$

from where we easily deduce the effective definition given in (II.39) and the number c_h.

[1] Z. Bern, S. Davies, T. Dennen and Y. -t. Huang, "Absence of Three-Loop Four-Point Divergences in $\mathcal{N} = 4$ Supergravity," Phys. Rev. Lett. **108** (2012) 201301 [arXiv:1202.3423 [hep-th]].

[2] P. Tourkine and P. Vanhove, "An R^4 Non-Renormalisation Theorem in $\mathcal{N} = 4$ Supergravity," Class. Quant. Grav. **29** (2012) 115006 [arXiv:1202.3692 [hep-th]].

[3] G. Bossard, P. S. Howe, K. S. Stelle and P. Vanhove, "The Vanishing Volume of $d = 4$ Superspace," Class. Quant. Grav. **28** (2011) 215005 [arXiv:1105.6087 [hep-th]].

[4] S. Deser, J. H. Kay and K. S. Stelle, "Renormalizability Properties of Supergravity," Phys. Rev. Lett. **38** (1977) 527.

[5] R. Kallosh, "On Absence of 3-Loop Divergence in $\mathcal{N} = 4$ Supergravity," Phys. Rev. D **85** (2012) 081702 [arXiv:1202.4690 [hep-th]].

[6] Z. Bern, J. J. Carrasco, L. J. Dixon, H. Johansson, D. A. Kosower and R. Roiban, “Three-Loop Superfiniteness of $\mathcal{N} = 8$ Supergravity,” Phys. Rev. Lett. **98** (2007) 161303 [hep-th/0702112].

[7] Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson and R. Roiban, “Manifest Ultraviolet Behavior for the Three-Loop Four-Point Amplitude of $\mathcal{N} = 8$ Supergravity,” Phys. Rev. D **78** (2008) 105019 [arXiv:0808.4112 [hep-th]].

[8] Z. Bern, J. J. Carrasco, L. J. Dixon, H. Johansson and R. Roiban, “The Ultraviolet Behavior of $\mathcal{N} = 8$ Supergravity at Four Loops,” Phys. Rev. Lett. **103** (2009) 081301 [arXiv:0905.2326 [hep-th]].

[9] M. B. Green, J. G. Russo and P. Vanhove, “Non-Renormalisation Conditions in Type II String Theory and Maximal Supergravity,” JHEP **0702** (2007) 099 [hep-th/0610299].

[10] M. B. Green, J. G. Russo and P. Vanhove, “String Theory Dualities and Supergravity Divergences,” JHEP **1006** (2010) 075 [arXiv:1002.3805 [hep-th]].

[11] G. Bossard, P. S. Howe and K. S. Stelle, “The Ultra-Violet Question in Maximally Supersymmetric Field Theories,” Gen. Rel. Grav. **41** (2009) 919 [arXiv:0901.4661 [hep-th]].

[12] G. Bossard, P. S. Howe and K. S. Stelle, “On Duality Symmetries of Supergravity Invariants,” JHEP **1101** (2011) 020 [arXiv:1009.0743 [hep-th]].

[13] A. A. Tseytlin, “Heterotic type I superstring duality and low-energy effective actions,” Nucl. Phys. B **467** (1996) 383 [hep-th/9512081].

[14] C. Bachas, C. Fabre, E. Kiritsis, N. A. Obers and P. Vanhove, “Heterotic / Type I Duality and D-Brane Instantons,” Nucl. Phys. B **509** (1998) 33 [hep-th/9707126].

[15] E. D’Hoker and D. H. Phong, “Two-Loop Superstrings Vi: Non-Renormalization Theorems and the 4-Point Function,” Nucl. Phys. B **715** (2005) 3 [hep-th/0501197].

[16] H. Elvang, D. Z. Freedman and M. Kiermaier, “A Simple Approach to Counterterms in $\mathcal{N} = 8$ Supergravity,” JHEP **1011** (2010) 016 [arXiv:1003.5018 [hep-th]].

[17] H. Elvang and M. Kiermaier, “Stringy Klt Relations, Global Symmetries, and E7(7) Violation,” JHEP **1010** (2010) 108 [arXiv:1007.4813 [hep-th]].

[18] N. Beisert, H. Elvang, D. Z. Freedman, M. Kiermaier, A. Morales and S. Stieberger, “E7(7) Constraints on Counterterms in $\mathcal{N} = 8$ Supergravity,” Phys. Lett. B **694** (2010) 265 [arXiv:1009.1643 [hep-th]].

[19] H. Elvang, D. Z. Freedman and M. Kiermaier, “SUSY Ward Identities, Superamplitudes, and Counterterms,” J. Phys. A A **44** (2011) 454009 [arXiv:1012.3401 [hep-th]].

[20] A. Dabholkar and J. A. Harvey, "String Islands," JHEP **9902** (1999) 006 [hep-th/9809122].

[21] E. Witten, "String Theory Dynamics in Various Dimensions," Nucl. Phys. B **443** (1995) 85 [arXiv:hep-th/9503124].

[22] C. M. Hull and P. K. Townsend, "Unity of Superstring Dualities," Nucl. Phys. B **438** (1995) 109 [arXiv:hep-th/9410167].

[23] A. Gregori, E. Kiritsis, C. Kounnas, N. A. Obers, P. M. Petropoulos and B. Pioline, "R^2 Corrections and Non-Perturbative Dualities of $\mathcal{N} = 4$ String Ground States," Nucl. Phys. B **510** (1998) 423 [arXiv:hep-th/9708062].

[24] D. C. Dunbar and P. S. Norridge, "Calculation of Graviton Scattering Amplitudes Using String Based Methods," Nucl. Phys. B **433** (1995) 181 [hep-th/9408014].

[25] D. C. Dunbar and P. S. Norridge, "Infinities Within Graviton Scattering Amplitudes," Class. Quant. Grav. **14** (1997) 351 [arXiv:hep-th/9512084].

[26] D. C. Dunbar, J. H. Ettle and W. B. Perkins, "Perturbative Expansion of N¡8 Supergravity," Phys. Rev. D **83** (2011) 065015 [arXiv:1011.5378 [hep-th]].

[27] D. C. Dunbar, J. H. Ettle and W. B. Perkins, "Obtaining One-Loop Gravity Amplitudes Using Spurious Singularities," Phys. Rev. D **84** (2011) 125029 [arXiv:1109.4827 [hep-th]].

[28] D. C. Dunbar, J. H. Ettle and W. B. Perkins, "The N-Point Mhv One-Loop Amplitude in $\mathcal{N} = 4$ Supergravity," Phys. Rev. Lett. **108** (2012) 061603 [arXiv:1111.1153 [hep-th]].

[29] D. C. Dunbar, J. H. Ettle and W. B. Perkin, "Constructing Gravity Amplitudes from Real Soft and Collinear Factorisation," arXiv:1203.0198 [hep-th].

[30] Z. Bern, C. Boucher-Veronneau and H. Johansson, "$N \geq 4$ Supergravity Amplitudes from Gauge Theory at One Loop," Phys. Rev. D **84** (2011) 105035 [arXiv:1107.1935 [hep-th]].

[31] S. Ferrara and C. Kounnas, "Extended Supersymmetry in Four-Dimensional Type Ii Strings," Nucl. Phys. B **328** (1989) 406.

[32] S. Chaudhuri and J. Polchinski, "Moduli Space of Chl Strings," Phys. Rev. D **52** (1995) 7168 [arXiv:hep-th/9506048].

[33] J. H. Schwarz and A. Sen, "Type Iia Dual of the Six-Dimensional Chl Compactification," Phys. Lett. B **357** (1995) 323 [arXiv:hep-th/9507027].

[34] P. S. Aspinwall, "Some Relationships Between Dualities in String Theory," Nucl. Phys. Proc. Suppl. **46** (1996) 30 [arXiv:hep-th/9508154].

[35] N. Sakai and Y. Tanii, "One Loop Amplitudes and Effective Action in Superstring Theories,"

Nucl. Phys. B **287** (1987) 457.

[36] K. S. Narain, M. H. Sarmadi and E. Witten, “A Note on Toroidal Compactification of Heterotic String Theory,” Nucl. Phys. B **279** (1987) 369.

[37] D. P. Jatkar and A. Sen, “Dyon Spectrum in Chl Models,” JHEP **0604** (2006) 018 [arXiv:hep-th/0510147].

[38] A. Dabholkar and D. Gaiotto, “Spectrum of Chl Dyons from Genus-Two Partition Function,” JHEP **0712** (2007) 087 [arXiv:hep-th/0612011].

[39] S. ’y. Mizoguchi, “On Asymmetric Orbifolds and the D = 5 No-Modulus Supergravity,” Phys. Lett. B **523** (2001) 351 [hep-th/0109193].

[40] A. Sen and C. Vafa, “Dual Pairs of Type II String Compactification,” Nucl. Phys. B **455** (1995) 165 [arXiv:hep-th/9508064].

[41] M. B. Green and P. Vanhove, “The Low Energy Expansion of the One-Loop Type II Superstring Amplitude,” Phys. Rev. D **61** (2000) 104011 [arXiv:hep-th/9910056].

[42] N. E. J. Bjerrum-Bohr and P. Vanhove, “Absence of Triangles in Maximal Supergravity Amplitudes,” JHEP **0810** (2008) 006 [arXiv:0805.3682 [hep-th]].

[43] M. T. Grisaru and W. Siegel, “The One Loop Four Particle S Matrix in Extended Supergravity,” Phys. Lett. B **110** (1982) 49.

[44] Z. Bern, J. J. Carrasco, D. Forde, H. Ita and H. Johansson, “Unexpected Cancellations in Gravity Theories,” Phys. Rev. D **77** (2008) 025010 [arXiv:0707.1035 [hep-th]].

[45] G. ’t Hooft and M. J. G. Veltman, “One Loop Divergencies in the Theory of Gravitation,” Annales Poincare Phys. Theor. A **20** (1974) 69.

[46] M. B. Green, J. H. Schwarz and L. Brink, “$\mathcal{N} = 4$ Yang-Mills and $\mathcal{N} = 8$ Supergravity as Limits of String Theories,” Nucl. Phys. B **198** (1982) 474.

[47] M. Fischler, “Finiteness Calculations for $O(4)$ Through $O(8)$ Extended Supergravity and $O(4)$ Supergravity Coupled to Selfdual $O(4)$ Matter,” Phys. Rev. D **20** (1979) 396.

[48] I.S. Gradshteyn, and I.W Ryzhik, *Table of Integrals*, Series, and Products (New York: Academic Press), 1980